

Recombinant Human VTCN1 protein, Fc-tagged, low endotoxin

Cat. No. VTCN1-572H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human VTCN1 (Phe29-Ala258) protein (low endotoxin) was fused to human IgG1 Fc tag at C-terminus and expressed in human 293 cells (HEK293).
Species	Human
Source	HEK293
ProteinLength	Phe29-Ala258
Description	V-set domain-containing T-cell activation inhibitor 1 (VTCN1) is also known as Immune costimulatory protein B7-H4, Protein B7S1, T-cell costimulatory molecule B7x, B7H4, which belongs to the immunoglobulin superfamily and BTN/MOG family. VTCN1 contains two Ig-like V-type (immunoglobulin-like) domains. The expression of VTCN1 is up-regulated by IL6 and IL10 and is inhibited by GM-CSF and IL4 on antigen-presenting cells (APCs). VTCN1/B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation.
Predicted N Terminal	Phe29
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 51.8 kDa. The protein migrates as 66-85 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	VTCN1
Official Symbol	VTCN1
Synonyms	VTCN1; V-set domain containing T cell activation inhibitor 1; V-set domain-containing T-cell activation inhibitor 1; B7 family member; H4; B7 superfamily member 1; B7 H4; B7H4; B7S1; B7X; FLJ22418; B7 family member, H4; T cell costimulatory molecule B7x; T-cell costimulatory molecule B7x; immune costimulatory protein B7-H4; B7-H4; B7h.5; VCTN1; PRO1291; RP11-229A19.4
Gene ID	79679
mRNA Refseq	NM_001253849
Protein Refseq	NP_001240778

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM 608162


UniProt ID Q7Z7D3

**Bioactivity-ELISA of
VTCN1-572H**



Immobilized Human B7-H4, Fc Tag, low endotoxin at 0.5 µg/mL (100 µL/well) can bind Anti-B7-H4 MAb (Human IgG1) with a linear range of 0.2-8 ng/mL (QC tested).

**SDS-PAGE of
VTCN1-572H**



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Human B7-H4, Fc Tag, low endotoxin on SDS-PAGE under reducing (R) condition.
The gel was stained overnight with Coomassie Blue. The purity of the protein is
greater than 95%.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA