

Recombinant Mouse ICOS protein (C137S, C138S), Fc-tagged

Cat. No. ICOS-729M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse ICOS (C137S (Glu21-Leu144) protein was fused to human IgG1 Fc tag at C-terminus and expressed in human 293 cells (HEK293).
Species	Mouse
Source	HEK293
ProteinLength	Glu21-Leu144
Description	Inducible T-cell costimulator (ICOS) is also known as Activation-inducible lymphocyte immunomediatory molecule (AILIM), CD278, which belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4 and PD-1. ICOS enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. CD278/ICOS prevents the apoptosis of pre-activated T-cells and also plays a critical role in CD40-mediated class switching of immunoglobulin isotypes.
Predicted N Terminal	Glu21
Form	Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 40.6 kDa. The protein migrates as 45-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION


Gene Name	ICOS
Official Symbol	ICOS
Synonyms	ICOS; inducible T cell co-stimulator; inducible T-cell costimulator; CD28-related protein 1; CD28 and CTLA-4-like protein; inducible T-cell co-stimulator; activation-inducible lymphocyte immunomediatory molecule; H4; CCLP; AILIM; CRP-1; Ly115
Gene ID	54167
mRNA Refseq	NM_017480
Protein Refseq	NP_059508
UniProt ID	Q9WVS0

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity-ELISA of ICOS-729M



Immobilized Human B7-H2, His Tag at 1 µg/mL (100 µL/well) can bind Mouse ICOS (C137S, C138S), Fc Tag with a linear range of 0.3-10 ng/mL (QC tested).

SDS-PAGE of ICOS- 729M



Mouse ICOS (C137S, C138S), Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.