

Recombinant COVID-19 S protein RBD (R408I), His-tagged

Cat. No. Spike-216V **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant COVID-19 S protein RBD (R408I, Arg319-Phe541) was fused to His-tag at the C-terminus and expressed in human 293 cells (HEK293).
Species	Sars-CoV-2
Source	HEK293
ProteinLength	Arg319-Phe541
Description	It's been reported that COVID-19 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.
Predicted N Terminal	Arg319
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 27.0 kDa. The protein migrates as 33-35 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU per µg by the LAL method.


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
GENE INFORMATION	
Gene Name	S
Official Symbol	S
Synonyms	spike glycoprotein
Gene ID	43740568
Protein Refseq	YP_009724390.1
UniProt ID	P0DTC2

Bioactivity-ELISA of Spike-216V




Immobilized Human ACE2, Fc Tag at 2 µg/mL (100 µL/well) can bind COVID-19 S protein RBD (R408I), His Tag with a linear range of 0.5-8 ng/mL (QC tested).

Bioactivity-ELISA of Spike-216V




Immobilized COVID-19 S protein RBD (R408I), His Tag at 2 µg/mL (100 µL/well) can bind Monoclonal Anti-COVID-19-S protein RBD Antibody, Human IgG1 with a linear range of 0.2-3 ng/mL (Routinely tested).

Bioactivity-ELISA of Spike-216V




Immobilized COVID-19 S protein RBD (R408I), His Tag at 2 µg/mL (100 µL/well) can bind Anti-COVID-19 Neutralizing Antibody, Human IgG1 with a linear range of 0.195-6.25 ng/mL (Routinely tested).

Bioactivity-SPR of Spike-216V




Human ACE2, Fc Tag captured on CM5 chip via anti-human IgG Fc antibodies surface can bind COVID-19 S protein RBD (R408I), His Tag with an affinity constant of 5.35 nM as determined in a SPR assay (Biacore T200) (Routinely tested).

Bioactivity-BLI of Spike-216V



Loaded Human ACE2, Fc Tag on Protein A Biosensor, can bind COVID-19 S protein RBD (R408I), His Tag with an affinity constant of 4.47 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

SDS-PAGE of Spike- 216V



COVID-19 S protein RBD (R408I), His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.