

Active Recombinant Cupriavidus necator tdh Protein, His-tagged

Cat. No. tdh-01C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Cupriavidus necator tdh Protein with His tag was expressed in E.coli.
Species	Cupriavidus necator
Source	E.coli
Description	Threonine dehydrogenase (EC 1.1.1.103) is a β -NAD ⁺ coenzyme dependent enzyme, from the group of short chain alcohol dehydrogenases. It is a key enzyme in L threonine catabolism in microorganisms and mammals. The enzyme catalyzes conversion of l-threonine to 2-amino-3 ketobutyrate, with simultaneous reduction of NAD ⁺ .
Form	Lyophilized
Bio-activity	Specific activity is ≥ 2 U/mg
Molecular Mass	36.8 kDa
Purity	$\geq 95\%$
Usage	For Research Use Only! Not For Use in Humans.
Storage	Store the lyophilized protein at -20 centigrade. Reconstituted enzyme can be stored in working aliquots at -20 centigrade. Avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage Buffer Proprietary buffer

Reconstitution Centrifuge the vial prior to opening. Reconstitute in distilled water.

GENE INFORMATION

Synonyms ThrDH; tdh; L-threonine 3-dehydrogenase; Threonine Dehydrogenase

UniProt ID [E5RQ20](#)

Activity of Active ThrDH

Specific Activity of Threonine Dehydrogenase is ≥ 2 U/mg. In this reaction 1 μ g of ThrDH converts L-Threonine into 2-amino-3-oxobutyrates with concomitant reduction of NAD⁺ which was measured at 340 nm.

Size exclusion chromatography of ThrDH

SEC analysis of Threonine Dehydrogenase using a Superose 6 Increase 5x150 column 50 mM sodium phosphate; 0.3 M NaCl pH 7.2 at 100 μ min

SDS-PAGE (4-20%) of recombinant ThrDH

Recombinant Protein loaded under reducing conditions and stained with Coomassie Blue.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA