

Recombinant Human Nodal

Cat. No. NODAL-30H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human Nodal Protein, Tag Free, was expressed in E. coli.
Species	Human
Source	E.coli
Protein Length	His238-Leu347aa
Description	Nodal is a 13 kDa member of the TGF-beta superfamily of molecules (1-3). Its name derives from the fact that it is expressed in the primitive node of the developing embryo. In human, it is synthesized as a 347 amino acid (aa) preproprecursor that contains a 26 aa signal sequence, a 211 aa prodomain, and a 110 aa mature region (4, 5). Consistent with its TGF-beta superfamily membership, it exists as a disulfide-linked homodimer and would be expected to demonstrate a cysteine-knot motif (1, 5, 6). Mature human Nodal is 99%, 98%, 96% and 98% aa identical to mature canine, rat, bovine and mouse Nodal, respectively. Nodal signals through two receptor complexes, both of which contain members of the TGF-beta family of Ser/Thr kinase receptors. The first receptor complex is composed of ActRIB/ALK4 and ActRIIB. Nodal interaction with the ligand-binding subunit ALK4 requires the participation of fucosylated Cripto, a member of the EGF-CFC gene family. The second complex is composed of ALK7 and ActRIIB, and does not require Cripto interaction (7-9). Nodal signaling is regulated by a number of inhibitors. Nodal is reported to form nonsignaling disulfide-linked heterodimers with BMP-3, -4 and -7 (2, 6). There are also at least three secreted factors that block Nodal activation of its receptor. The first

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

is Cerberus, a member of the DAN family of proteins. The last two are TGF-beta family members named Lefty-A and B (2, 11). The exact mechanisms employed are unclear, but Leftys are apparently induced by nodal and can work at a considerable distance from their origin (2). Nodal is known to induce both mesoderm and endoderm, participate in anterior-posterior positioning, and through ALK7, induce apoptosis in susceptible cell populations (2, 8, 12).

Tag	Non
Bio-activity	Measured by its ability to induce Smad2 phosphorylation in P19 mouse embryonal carcinoma cells. Yeo, C. et al. (2001) Mol. Cell. 7:949. Approximately 100 ng/mL of Recombinant Human Nodal can effectively induce Smad2 phosphorylation.
Molecular Mass	12.9 kDa
Purity	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.
N-terminal Sequence Analysis	Met
Stability	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from a 0.2 µm filtered solution in Acetonitrile and TFA with BSA as a

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

carrier protein.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Reconstitution

Reconstitute at 100 µg/mL in sterile 4 mM HCl containing at least 0.1% human or bovine serum albumin.

GENE INFORMATION**Gene Name**

NODAL nodal growth differentiation factor [Homo sapiens (human)]

Official Symbol

NODAL

Synonyms

NODAL

Gene ID

4838

UniProt ID

Q96S42

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA