

Recombinant Mouse Camk2d Protein, DDK-tagged

Cat. No. Camk2d-030M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Purified recombinant protein of Mouse calcium/calmodulin-dependent protein kinase II, delta (Camk2d), transcript variant 2, mutant(T287A), with C-terminal DDK tag, expressed in sf9 cell.
Species	Mouse
Source	Insect Cells
Description	Calcium/calmodulin-dependent protein kinase involved in the regulation of Ca²⁺ homeostasis and excitation-contraction coupling (ECC) in heart by targeting ion channels, transporters and accessory proteins involved in Ca²⁺ influx into the myocyte, Ca²⁺ release from the sarcoplasmic reticulum (SR), SR Ca²⁺ uptake and Na⁺ and K⁺ channel transport. Targets also transcription factors and signaling molecules to regulate heart function. In its activated form, is involved in the pathogenesis of dilated cardiomyopathy and heart failure. Contributes to cardiac decompensation and heart failure by regulating SR Ca²⁺ release via direct phosphorylation of RYR2 Ca²⁺ channel on 'Ser-2808'. In the nucleus, phosphorylates the MEF2 repressor HDAC4, promoting its nuclear export and binding to 14-3-3 protein, and expression of MEF2 and genes involved in the hypertrophic program. Is essential for left ventricular remodeling responses to myocardial infarction. In pathological myocardial remodeling acts downstream of the beta adrenergic receptor signaling cascade to regulate key proteins involved in ECC. Regulates Ca²⁺ influx to myocytes by binding and phosphorylating the L-type Ca²⁺ channel subunit beta-2 CACNB2. In addition to Ca²⁺ channels, can target and regulate the cardiac

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

sarcolemmal Na⁺ channel Nav1.5/SCN5A and the K⁺ channel Kv4.3/KCND3, which contribute to arrhythmogenesis in heart failure. Phosphorylates phospholamban (PLN/PLB), an endogenous inhibitor of SERCA2A/ATP2A2, contributing to the enhancement of SR Ca²⁺ uptake that may be important in frequency-dependent acceleration of relaxation (FDAR) and maintenance of contractile function during acidosis. May participate in the modulation of skeletal muscle function in response to exercise, by regulating SR Ca²⁺ transport through phosphorylation of PLN/PLB and triadin, a ryanodine receptor-coupling factor.

Molecular Mass	55.3 kDa
Purity	>80%, as determined by SDS-PAGE and Coomassie blue staining
Stability	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Storage	Store at -80 centigrade.
Concentration	>50 ug/mL as determined by microplate BCA method
Storage Buffer	50mM Tris-HCl, pH8.0, 100mM glycine, 10% glycerol.

GENE INFORMATION

Gene Name	Camk2d calcium/calmodulin-dependent protein kinase II, delta [Mus musculus (house mouse)]
Official Symbol	Camk2d
Synonyms	Camk2d; calcium/calmodulin-dependent protein kinase II, delta; CaMK II; [d]-CaMKII; 2810011D23Rik; 8030469K03Rik; calcium/calmodulin-dependent protein kinase type

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


II subunit delta; caM kinase II subunit delta; caMK-II subunit delta; EC 2.7.11.17

Gene ID 108058

mRNA Refseq NM_001025438

Protein Refseq NP_001020609

UniProt ID Q6PHZ2



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA