

Recombinant Human IGHG3 Protein (SPR verified)

Cat. No. IGHG3-928H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IGHG3 protein (SPR verified)(Glu99-Lys377) was expressed in human 293 cells (HEK293).
Species	Human
Source	HEK293
ProteinLength	Glu99-Lys377
Description	Immunoglobulin G3 (IgG3) is a member of many immunoglobulin G developed and secreted by effective B cells. In wake of cutting by pepsin, IgG is divided into two F(ab)s with one antigen binding site and a high conserved Fc segment. The Fc segment bears a highly conserved N-glycosylation site. Ig gamma-3 chain C region (IgG3-Fc / IGHG3) contains two constant regions of IgG3 H chain (CH2, CH3).
Predicted N Terminal	Glu99
Form	Tris with Glycine, Arginine and NaCl, pH7.5
Molecular Mass	The protein has a calculated MW of 31.2 kDa. The protein migrates as 35-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

-20°C to -70°C for 12 months in lyophilized state;

-70°C for 3 months under sterile conditions after reconstitution.

Reconstitution

It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 mg/ml. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION
Gene Name

IGHG3

Official Symbol

IGHG3

Synonyms

Ig gamma-3 chain C region,IgG,IgG3

Gene ID


3502

MIM

147120


UniProt ID

P01860

Bioactivity-SPR of IGHG3-928H


Human FCGRT&B2M Heterodimer Protein, His Tag (SPR & BLI verified) captured on CM5 Chip via anti-His antibody can bind Human IgG3 Fc, Tag Free (SPR verified) with an affinity constant of 0.683 μ M as determined in SPR assay (Biacore 8K) (QC tested).

SDS-PAGE of IGHG3-928H



Human IgG3 Fc, Tag Free (SPR verified) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.