

Mouse Anti-IL8 Monoclonal Antibody

Cat. No. DMABT-H19230 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse Anti-IL8 Monoclonal Antibody
Species	Human
Source	Mouse
Antigen Description	IL-8, also known as neutrophil chemotactic factor, neutrophil activating protein, and monocyte-derived neutrophil chemotactic factor, is a member of the alpha (C-X-C) subfamily of chemokines called CXCL8. In response to proinflammatory stimuli, IL-8 is produced by monocytes, macrophages, T cells, neutrophils, and fibroblasts. IL-8 promotes neutrophil chemotaxis and degranulation. The 72 amino acid IL-8 is the predominant form secreted by monocytes and lymphocytes.
Target	IL8
Isotype	IgG1, kappa
species	Human
Clone	I9B6
Conjugation	N/A
Applications	ELISA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Preparation	The antibody was purified by affinity chromatography.
Usage recommendation	Each lot of this antibody is quality control tested by ELISA assay. For ELISA capture applications, a concentration range of 2-6 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of IL-8 recombinant protein ranging from 2000 to 26pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.
Concentration	0.5 mg/ml
Storage	The antibody solution should be stored undiluted at 4 °C.
Size	50 µg; 500 µg

GENE INFORMATION

Gene Name	IL8 interleukin 8 [Homo sapiens]
Official Symbol	IL8
Synonyms	IL8; interleukin 8; interleukin-8; 410C; alveolar macrophage chemotactic factor I; AMCF I; b ENAP; beta endothelial cell derived neutrophil activating peptide; chemokine (C X C motif) ligand 8; CXCL8; GCP 1; GCP1; granulocyte chemotactic protein 1; IL 8; K60; LECT; LUCT; lung giant cell carcinoma derived chemotactic protein; lymphocyte derived neutrophil activating peptide; LYNAP; MDNCF; MONAP; monocyte derived neutrophil chemotactic factor; monocyte derived neutrophil activating peptide; NAF; NAP 1; NAP1; neutrophil activating peptide 1; SCYB8; TSG 1; tumor necrosis factor induced gene 1; emoctakin; T-cell chemotactic factor; neutrophil-activating peptide 1; chemokine (C-X-C motif) ligand 8; beta-thromboglobulin-like protein; tumor necrosis factor-induced gene 1; monocyte-derived

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	neutrophil chemotactic factor; monocyte-derived neutrophil-activating peptide; small inducible cytokine subfamily B, member 8; lung giant cell carcinoma-derived chemotactic protein; beta endothelial cell-derived neutrophil activating peptide; GCP-1; NAP-1;
Gene ID	3576
mRNA Refseq	NM_000584
Protein Refseq	NP_000575
MIM	146930
UniProt ID	P10145
Chromosome Location	4q13-q21
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Activation of Genes by ATF4, organism-specific biosystem; Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem;
Function	chemokine activity; interleukin-8 receptor binding; protein binding;