

Recombinant Human NUP98 protein, hFC-tagged

Cat. No. NUP98-5793H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Recombinant Human NUP98 protein(NP_005378.4)(1-880 aa) was fused to hFC-tagged at C-terminus and expressed in HEK293.

Species

Human

Source

HEK293

ProteinLength

Partial

Description

Nuclear pore complexes (NPCs) regulate the transport of macromolecules between the nucleus and cytoplasm, and are composed of many polypeptide subunits, many of which belong to the nucleoporin family. This gene belongs to the nucleoporin gene family and encodes a 186 kDa precursor protein that undergoes autoproteolytic cleavage to generate a 98 kDa nucleoporin and 96 kDa nucleoporin. The 98 kDa nucleoporin contains a Gly-Leu-Phe-Gly (GLGF) repeat domain and participates in many cellular processes, including nuclear import, nuclear export, mitotic progression, and regulation of gene expression. The 96 kDa nucleoporin is a scaffold component of the NPC. Proteolytic cleavage is important for targeting of the proteins to the NPC. Translocations between this gene and many other partner genes have been observed in different leukemias. Rearrangements typically result in chimeras with the N-terminal GLGF domain of this gene to the C-terminus of the partner gene. Alternative splicing results in multiple transcript variants encoding different isoforms, at least two of which are proteolytically processed. Some variants lack the region that encodes the 96 kDa nucleoporin.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.
Molecular Mass	116 kDa
AA Sequence	MFNKSFGTPFGGGTGGFGTTSTFGQNTGFGTSSGGAFGTSAFGSSNNTGGFLGNS QTKPGGLFGTSSFSQPATSTSTGFGFGTSTGTANTLFGTASTGTSLFSSQNNAFAQ NKPTGFGNFGTSTSSGGLFGTTNTTNSPFGSTSGSLFGPSSFTAAPTGTTIKFNPTT GDTMVKAGVSTNISTKHQCITAMKEYESKSLEELRLEDYQANRKGPNQVAGATT TGLFGSSPATSSATGLFSSSTTNSGFAYGQNKTAFGTSTTGFGTNPGGFLGQQNQ QTTSLFSKPFQGATTTQNTGFSFGNTSTIGQPSTNTMGLFGVTQASQPGGLFGTAT NTSTGTAFGTGTGLFGQNTGFGAVGSTLFGNNKLTTFGSSTTSAPSFGTSSGGLF GNKPTLTLGTNTNTSNFGFGTNTSGNSIFGSKPAPGTLGTGLGAGFGTALGAGQAS LFGNNQPKIGGPLGTGAFGAPGFNTTTATLGFGAPQAPVALTDPNASAAQQAVLQQ HINSLTYSPFGDSPLFRNPMSDPKKKEERLKPTNPAAQKALTTPHYKLTPRPATRV RPKALQTTGTAKSHLFDGLDDDEPSLANGAFMPKKSIIKLVLKLNLSNLFSPVNRD SENLASPSEYPENGERFSFLSKPVDENHQDGDDESLVSHFYTNPIAKPIQPESAS GNKHSNSNSVDDTIVALNMRAALRNGLEGSSEETSFHDESLQDDREEIENNSYHMH PAGIILTKVGYYTIPSMDDLAKITNEKGECIVSDFTIGRKGYGSIYFEGDVNLNLNDD IVHIRRKEVVYLLDDNQKPPVGEGLNRKAEVTLDGWPTDKTSRCLIKSPDRLADINY EGRLEAVSRKQGAQFKEYRPETGSWVFKVSHF
Purity	Greater than 90% as determined by SDS-PAGE.
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Storage	Store at -20°C upon receipt, aliquoting is necessary for multiple use. Avoid repeated freeze-thaw cycles.

Reconstitution

It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 mg/ml. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION
Gene Name

NUP98

Official Symbol

NUP98

Synonyms

96 kDa nucleoporin; 98 kDa nucleoporin; ADAR2; ADIR2 ; GLFG-repeat containing nucleoporin; Nuclear pore complex protein Nup96; Nuclear pore complex protein Nup98 Nup96; Nucleoporin 98kD ; nucleoporin 98kDa; Nucleoporin Nup96; Nucleoporin Nup98; NUP196 ; NUP96; Nup98; Nup98-Nup96; NUP98_HUMAN

Gene ID

4928

mRNA Refseq

NM_005387.7

Protein Refseq

NP_005378.4

UniProt ID


P52948

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**SDS-PAGE of
NUP98-5793H**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA