

Recombinant Mouse Icam2 Protein, MYC/DDK-tagged

Cat. No. Icam2-1205M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Purified recombinant protein of full-length mouse intercellular adhesion molecule 2 (Icam2), with C-terminal MYC/DDK tag, was expressed in HEK293T cells.
Species	Mouse
Source	HEK293
Description	ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.
Molecular Mass	31.4 kDa
Purity	>80% as determined by SDS-PAGE and Coomassie blue staining
Stability	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Storage	Store at -80 centigrade after receiving vials.
Concentration	>50 g/mL as determined by microplate BCA method
Storage Buffer	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	Icam2 intercellular adhesion molecule 2 [Mus musculus (house mouse)]
Official Symbol	Icam2
Synonyms	Icam2; intercellular adhesion molecule 2; CD102; Ly-60; Icam-2; intercellular adhesion molecule 2; lymphocyte function-associated AG-1 counter-receptor
Gene ID	15896
mRNA Refseq	NM_010494
Protein Refseq	NP_034624
UniProt ID	P35330

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA