

Recombinant Human Mononucleosomes (H3.3 ΔN32)

Cat. No. NUC-010 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mononucleosomes assembled from recombinant human histones expressed in E. coli.
Species	Human
Source	E.coli
Description	Mononucleosomes assembled from recombinant human histones expressed in E. coli (two each of histones H2A, H2B, H3.3 and H4. Accession numbers: H2A-P04908; H2B-O60814; H3.3-P84243; H4-P62805), with the amino acid sequence of H3 beginning with glycine 33 (amino acids 1-32 are deleted), wrapped by 147 base pairs of 601 positioning sequence DNA. The nucleosome is the basic subunit of chromatin. The 601 sequence, identified by Lowary and Widom, is a 147-base pair sequence that has high affinity for histone octamers and is useful for nucleosome assembly.
Form	Mononucleosomes (H3.3 ΔN32), Recombinant Human (27.4 g protein weight, 50 g DNA+protein) in 10 mM Tris HCl, pH 7.5, 25 mM NaCl, 1 mM EDTA, 2 mM DTT, 20% glycerol.
Molecular Mass	193,766.7 Da
Applications	Mononucleosomes (H3.3 ΔN32), Human Recombinant are highly purified and suitable for use as substrates in enzyme screening assays, structural studies, or effector protein binding experiments. The N-terminal deletion allows for the study of the role of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

the N-terminus in many aspects of chromatin biology.

Storage

Stable for six months at -80°C from date of receipt. For best results, aliquot and avoid multiple freeze/thaws.

Protein Gel Data of NUC-010




Coomassie stained PAGE gel of proteins in Mononucleosomes (H3.3 ΔN32), Human Recombinant (1 g) demonstrates the purity of the histones in the preparation. Sizes of molecular weight markers and positions of the core histones (H2A, H2B, H3.3 ΔN32 and H4) are indicated. H3.3 ΔN32 and H4 co-migrate.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127


45-1 Ramsey Road, Shirley, NY 11967, USA

DNA Gel Data of NUC-010



Mononucleosomes (H3.3 Δ N32), Human Recombinant run on an agarose gel and stained with ethidium bromide to visualize DNA. Lane 1: DNA extracted from nucleosomes (100 ng). Lane 2: Intact nucleosomes (300 ng).

Western Blot Data of NUC-010



Western Analysis of Nucleosome (H3.3)

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Δ N32), Human Recombinant. Top Panel: Wild type (Lane 1) and H3.3 Δ N32-containing nucleosomes (Lane 2) were probed with an anti-H3 COOH-terminal antibody and analyzed via ECL readout.

Bottom Panel: Detail from Coomassie stained gel showing histones from H3.3 WT (Lane 1) and H3.3 Δ N32 nucleosomes (Lane 2).

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA