

Recombinant Human Nucleosome, H3K4me3, K9,14,18ac dNuc, Biotinylated

Cat. No. NUC-016 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mononucleosomes assembled from recombinant human histones expressed in E. coli.
Species	Human
Source	E.coli
Description	Mononucleosomes assembled from recombinant human histones expressed in E. coli (two each of histones H2A, H2B, H3 and H4; accession numbers: H2A-P04908; H2B-O60814; H3.2- Q71DI3 *; H4-P62805) wrapped by 147 base pairs of 601 positioning sequence DNA. Histone H3 (created by a proprietary semi-synthetic method) contains trimethyl-lysine at position 4, and acetyl-lysine at positions 9, 14, and 18. The nucleosome is the basic subunit of chromatin. The 601 sequence, identified by Lowary and Widom, is a 147-base pair sequence that has high affinity for histone octamers and is useful for nucleosome assembly and contains a 5' biotin-TEG group. * H3K4me3, K9,14,18ac also has a Cys to Ala substitution at position 110.
Form	Nucleosome, Recombinant Human, H3K4me3, K9,14,18ac (27.3 µg protein weight, 50 µg total weight) in 10 mM Tris HCl, pH 7.5, 25 mM NaCl, 1 mM EDTA, 2 mM DTT, 20% glycerol.
Molecular Mass	200,644 Da
Applications	Nucleosome, Recombinant Human, H3K4me3, K9,14,18ac dNucs are highly purified

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


and are suitable for use as substrates in enzyme screening assays or for effector protein binding experiments.

Storage

Stable for six months at -80°C from date of receipt. For best results, aliquot and avoid multiple freeze/thaws.

Conjugation

Biotin

**Western Blot Data of
NUC-016**


Western Analysis of Nucleosome, Recombinant Human, H3K4me3, K9,14,18ac.


Top Panel: Unmodified H3 (Lane 1) and H3K4me3, K9,14,18ac containing nucleosomes (Lanes 2-5) were probed with an anti-H3K4me3, anti-H3K9ac, anti-H3K14ac, and anti-H3K18ac antibodies and analyzed via ECL readout. Only the H3K4me3, K9,14,18ac sample produced a detectable signal. Bottom Panel: Detail from Coomassie stained gel showing unmodified nucleosomes (Lane 1) and H3K4me3, K9,14,18ac nucleosomes (Lanes 2-5).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Mass Spec Data of NUC-016



H3K4me3, K9,14,18ac protein analyzed by high resolution mass spectrometry.
Expected mass = 15,389.8 Da. Determined mass = 15,391.5 Da.

Protein Gel Data of NUC-016




Coomassie stained PAGE gel of proteins in Nucleosome, Recombinant Human, H3K4me3, K9,14,18ac (1 µg) to demonstrate the purity of the histones in the preparation. Sizes of molecular weight markers and positions of the core histones (H2A, H2B, H3K4me3, K9,14,18ac and H4) are indicated.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

**DNA Gel Data of
NUC-016**



Nucleosome, Recombinant Human, H3K4me3, K9,14,18ac run on a native PAGE gel and stained with ethidium bromide to visualize DNA. Lane 1: Free DNA (200 ng).
Lane 2: Intact nucleosomes (400 ng).

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA