

Recombinant Human Nucleosome, H2AXS139phos dNuc, Biotinylated

Cat. No. NUC-039 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mononucleosomes assembled from recombinant human histones expressed in E. coli.
Species	Human
Source	E.coli
Description	Mononucleosomes assembled from recombinant human histones expressed in E. coli (two each of histones H2AX, H2B, H3 and H4; accession numbers: H2AX-P16104; H2BO60814; H3.1-P68431; H4-P62805) wrapped by 147 base pairs of 601 positioning sequence DNA. Histone H2AX (created by a proprietary synthetic method) contains phosphorylated-serine at position 139. The nucleosome is the basic subunit of chromatin. The 601 sequence, identified by Lowary and Widom, is a 147-base pair sequence that has high affinity for histone octamers and is useful for nucleosome assembly and contains a 5' biotin-TEG group.
Form	Nucleosome, Recombinant Human, H2AXS139phos (27.4µg protein weight, 50 µg total weight) in µl of 10 mM Tris HCl, pH 7.5, 25 mM NaCl, 1 mM EDTA, 2 mM DTT, 20% glycerol.
Molecular Mass	202,066.8 Da
Applications	Nucleosome, Recombinant Human, H2AXS139phos dNucs are highly purified and are suitable for use as substrates in enzyme screening assays or for effector protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


binding experiments. Nucleosome, Recombinant Human, H2AXS139phos dNucs does not contain free DNA which could alter assayed activities.

Storage

Stable for six months at -80°C from date of receipt. For best results, aliquot and avoid multiple freeze/thaws.

Conjugation

Biotin

Western Blot Data



Western Analysis of Nucleosome, Recombinant Human, H2AXS139phos. Top Panel: Unmodified H2A (Lane 1) and H2AXS139phos containing nucleosomes (Lane 2) were probed with an anti-H2AXS139phos antibody and analyzed via ECL readout. Only the H2AXS139phos sample produced a detectable signal. Bottom Panel: Detail from Coomassie stained gel showing unmodified nucleosomes (Lane 1) and H2AXS139phos nucleosomes (Lane 2).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Mass Spec Data



H2AXS139phos protein analyzed by high resolution mass spectrometry. Expected mass = 15092.38 Da. Determined mass = 15,092.38 Da.

Protein Gel Data




Coomassie stained PAGE gel of proteins in Nucleosome, Recombinant Human, H2AXS139phos (1 µg) to demonstrate the purity of the histones in the preparation. Sizes of molecular weight markers and positions of the core histones (H2AXS139phos, H2B, H3 and H4) are indicated

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

DNA Gel Data



Nucleosome, Recombinant Human, H2AXS139phos run on a native PAGE gel and stained with ethidium bromide to visualize DNA. Lane 1: Free DNA (100 ng). Lane 2: Intact nucleosomes (400 ng).

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA