

Recombinant Human Mononucleosomes (H2AZ.1)

Cat. No. NUC-041 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mononucleosomes assembled from recombinant human histones expressed in E. coli.
Species	Human
Source	E.coli
Description	Mononucleosomes assembled from recombinant human histones expressed in E. coli (two each of histones H2AZ.1, H2B, H3 and H4. Accession numbers: H2AZ.1-P0C0S5; H2B-O60814; H3.1-P68431; H4-P62805) wrapped by 147 base pairs of 601 positioning sequence DNA. The nucleosome is the basic subunit of chromatin. H2AZ.1 and H2AZ.2 are variants of histone H2A and implicated in diverse cellular functions ranging from transcriptional regulation, chromosome transmission and DNA damage repair. H2AZ has a dedicated deposition machinery (the SWR-C ATPase).
Form	Purified recombinant mononucleosomes (50 g total mass, 27.4 g in 50.0 μl) in 10 mM Tris-HCl pH 7.5, 1 mM EDTA, 25 mM NaCl, 2 mM DTT, & 20% glycerol.
Molecular Mass	198,425.5 Da
Applications	Mononucleosomes (H2AZ.1), Human Recombinant are highly purified and are suitable for use as substrates in enzyme screening assays or for nucleosome binding experiments. The absence of post-translational histone modifications makes them ideal for conducting enzyme activity and screening assays. Mononucleosomes (H2AZ.1), Human Recombinant do not contain free DNA which could alter assayed

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

activities.

Storage

Stable for six months at -80°C from date of receipt. For best results, aliquot and avoid multiple freeze/thaws.

Protein Gel Data




Coomassie stained PAGE gel of proteins in Nucleosome (H2AZ.1), Human Recombinant (1 g) demonstrates the purity of the histones in the preparation. Sizes of molecular weight markers and positions of the core histones (H2AZ.1, H2B, H3 and H4) are indicated.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127


45-1 Ramsey Road, Shirley, NY 11967, USA

DNA Gel Data



Nucleosome (H2AZ.1), Human Recombinant resolved via native PAGE and stained with ethidium bromide to visualize DNA. Lane 1: DNA extracted from nucleosomes (100 ng). Lane 2: Intact nucleosomes (400 ng).

Western Blot Data



Western Analysis of Nucleosome (H2AZ.1), Human Recombinant. Top Panel: Wild

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

type (Lane 1) and H2AZ.1-containing nucleosomes (Lane 2) were probed with an anti-H2AZ antibody and analyzed via ECL readout. Only the H2AZ.1 samples produced a detectable signal. Bottom Panel: Detail from Coomassie stained gel showing histones from H2A (Lane 1) and H2AZ.1

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA