

Recombinant Human CD70 protein, His-tagged

Cat. No. CD70-196H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD70 protein(P32970-1)(Leu50-Pro193), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Leu50-Pro193
Tag	N-His
Form	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.
Bio-activity	Immobilized Human CD27 Ligand Trimer, His Tag at 2ug/ml (100ul/Well) on the plate. Dose response curve for Human CD27, hFc Tag with the EC50 of 13.5ng/ml determined by ELISA.
Molecular Mass	The protein has a predicted MW of 51.8 kDa. Due to glycosylation, the protein migrates to 62-65 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE;> 95% as determined by HPLC


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Reconstituted protein stable at -80°C for 12 months, 4°C for 1 week. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
GENE INFORMATION	
Gene Name	CD70 CD70 molecule [Homo sapiens]
Official Symbol	CD70
Synonyms	CD70; CD70 molecule; CD27LG, TNFSF7, tumor necrosis factor (ligand) superfamily, member 7; CD70 antigen; CD27L; CD27-L; CD27 ligand; Ki-24 antigen; surface antigen CD70; tumor necrosis factor ligand superfamily member 7; tumor necrosis factor (ligand) superfamily, member 7; CD27LG; TNFSF7;
Gene ID	970
mRNA Refseq	NM_001252
Protein Refseq	NP_001243
MIM	602840
UniProt ID	P32970

Tris-Bis PAGE



ELISA Data



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA