

Recombinant Human CD70 protein, His-Flag-tagged

Cat. No. CD70-089H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD70 Protein is expressed from mammalian with His and Flag at the N-terminal. It contains Gln50-Pro193 Trimer.
Species	Human
Source	HEK293
ProteinLength	It contains Gln50-Pro193 Trimer.
Description	CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.
Form	Supplied as 0.22um filtered solution in PBS (pH 7.4).
Molecular Mass	The protein has a calculated MW of 51.8kDa. As a result of glycosylation, the protein migrates to 62- 65kDa by Bis-Tris PAGE.
Endotoxin	Less than 0.001EU per ug by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE; > 95% as determined by HPLC
Storage	The product should be stored at -70°C. Please do not repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Reconstitution

It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name CD70 CD70 molecule [Homo sapiens]

Official Symbol CD70

Synonyms CD70 molecule; CD70; TNFSF7; TNFSF7CD70 antigen; CD27 Ligand; CD27-L; CD27LCD27 ligand; CD27LG; Ki-24 antigen; surface antigen CD70; tumor necrosis factor (ligand) superfamily, member 7; Tumor necrosis factor ligand superfamily member 7

Gene ID 970

mRNA Refseq NM_001252

Protein Refseq NP_001243

MIM 602840

UniProt ID P32970

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA