

Recombinant Human HSD11B2 Protein, Myc/DDK-tagged, C13 and N15-labeled

Cat. No. HSD11B2-5346H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	HSD11B2 MS Standard C13 and N15-labeled recombinant protein (NP_000187) with a C-terminal MYC/DDK tag, was expressed in HEK293 cells.
Species	Human
Source	HEK293
Description	There are at least two isozymes of the corticosteroid 11-beta-dehydrogenase, a microsomal enzyme complex responsible for the interconversion of cortisol and cortisone. The type I isozyme has both 11-beta-dehydrogenase (cortisol to cortisone) and 11-oxoreductase (cortisone to cortisol) activities. The type II isozyme, encoded by this gene, has only 11-beta-dehydrogenase activity. In aldosterone-selective epithelial tissues such as the kidney, the type II isozyme catalyzes the glucocorticoid cortisol to the inactive metabolite cortisone, thus preventing illicit activation of the mineralocorticoid receptor. In tissues that do not express the mineralocorticoid receptor, such as the placenta and testis, it protects cells from the growth-inhibiting and/or pro-apoptotic effects of cortisol, particularly during embryonic development. Mutations in this gene cause the syndrome of apparent mineralocorticoid excess and hypertension.
Molecular Mass	44.1 kDa
AA Sequence	MERWPWPSGGAWLLVAARALLQLLRSDLRLGRPLLAALALLAALDWLCQRLPPPA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ALAVLAAAGWIALSRLARPQRLPVATRAVLITGCDSGFGKETAKKLDMSGFTVLATVL
ELNSPGAIELRTCCSPRLRLLQMDLTKPGDISRVLEFTKAHTTSTGLWGLVNNAGHN
EVDADAELSPVATFRSCMEVNFFGALELTGKLLPLLSSRGRIVTVGSPAGDMPYPC
LGAYGTSKAAVALLMDTFSCCELLPWGVKVSIIQPGCFKTESVRNVGQWEKRKQLLLA
NLPQELLQAYGKDYIEHLHGQFLHSLRLAMSDLTPVVDAITDALLAARPRRRYYPGQ
GLGLMYFIHYLLPEGLRRRFLQAFFISHCLPRALQPGQPGTTPPQDAAQGNLSPGP
SPAVARTRTRPLEQKLISEEDLAANDILDYKDDDDKV

Purity > 80% as determined by SDS-PAGE and Coomassie blue staining

Stability Stable for 3 months from receipt of products under proper storage and handling conditions.

Storage Store at -80 centigrade. Avoid repeated freeze-thaw cycles.

Concentration 50 µg/mL as determined by BCA

Storage Buffer 100 mM glycine, 25 mM Tris-HCl, pH 7.3.

GENE INFORMATION

Gene Name HSD11B2 hydroxysteroid 11-beta dehydrogenase 2 [Homo sapiens (human)]

Official Symbol HSD11B2

Synonyms HSD11B2; hydroxysteroid (11-beta) dehydrogenase 2; corticosteroid 11-beta-dehydrogenase isozyme 2; SDR9C3; short chain dehydrogenase/reductase family 9C; member 3; 11-DH2; 11-beta-HSD2; -HSD11 type II; 11-beta-hydroxysteroid dehydrogenase type 2; 11-beta-hydroxysteroid dehydrogenase type II; NAD-dependent 11-beta-hydroxysteroid dehydrogenase; short chain dehydrogenase/reductase family 9C member 3; AME; AME1; HSD2; HSD11K;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	3291
mRNA Refseq	NM_000196
Protein Refseq	NP_000187
MIM	614232
UniProt ID	P80365
SDS-PAGE	