

Recombinant Human NARF Protein, Myc/DDK-tagged, C13 and N15-labeled

Cat. No. NARF-6610H Lot. No. (See product label)

SPECIFICATION

Product Overview	NARF MS Standard C13 and N15-labeled recombinant protein (NP_036468) with a C-terminal MYC/DDK tag, was expressed in HEK293 cells.
Species	Human
Source	HEK293
Description	Several proteins have been found to be prenylated and methylated at their carboxyl-terminal ends. Prenylation was initially believed to be important only for membrane attachment. However, another role for prenylation appears to be its importance in protein-protein interactions. The only nuclear proteins known to be prenylated in mammalian cells are prelamin A- and B-type lamins. Prelamin A is farnesylated and carboxymethylated on the cysteine residue of a carboxyl-terminal CaaX motif. This post-translationally modified cysteine residue is removed from prelamin A when it is endoproteolytically processed into mature lamin A. The protein encoded by this gene binds to the prenylated prelamin A carboxyl-terminal tail domain. It may be a component of a prelamin A endoprotease complex. The encoded protein is located in the nucleus, where it partially colocalizes with the nuclear lamina. It shares limited sequence similarity with iron-only bacterial hydrogenases. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene, including one with a novel exon that is generated by RNA editing.
Molecular Mass	51 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MKCEHCTRKECSKKTCTDDQENV SADAPSPAQENGEKGEFHKLADAKIFLSDCLAC DSCMTAEEGVQLSQQNAKDFFRVLNLNKKCDTSKHKVLVSVCPQSLPYFAAKFNL SVTDASRRLCGFLKSLGVHYVFDTTIAADFSILESQKEFVRRYRQHSEEERTLPMLTS ACPGWVRYAERVLGRPITAHLC TAKSPQQVMGSLVKDYFARQQNLSPEKIFHVIVAP CYDKKLEALQESLPPALHGSRGADCVLTSGEIAQIMEQGDLSVRDAAVDTLFGDLKE DKVTRHDGASSDGH LAHIFRHA AKELFNEDVEEV TYRALRNKDFQEVTLEKNGEVV LRFAAAYGFRNIQNMILK LKKGKFPFHFVEVLACAGGCLNGRGQAQTPDGHADKAL LRQMEGIYADIPVRRPESSAHVQELYQE WLEGINS PKAREVLHTTYQS QERGTHSL DIKWSGPTRTRRLEQKLISEEDLAANDILDYKDDDDKV
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Stability	Stable for 3 months from receipt of products under proper storage and handling conditions.
Storage	Store at -80 centigrade. Avoid repeated freeze-thaw cycles.
Concentration	50 µg/mL as determined by BCA
Storage Buffer	100 mM glycine, 25 mM Tris-HCl, pH 7.3.
GENE INFORMATION	
Gene Name	NARF nuclear prelamins A recognition factor [Homo sapiens (human)]
Official Symbol	NARF
Synonyms	NARF; nuclear prelamins A recognition factor; DKFZp434G0420; FLJ10067; IOP2; iron only hydrogenase like protein 2; iron-only hydrogenase-like protein 2; prenyl-dependent prelamins A binding protein;

Gene ID	26502
mRNA Refseq	NM_012336
Protein Refseq	NP_036468
MIM	605349
UniProt ID	Q9UHQ1
SDS-PAGE	