

Recombinant Human MMP3 (Stromelysin 1, Progelatinase), Inactive, Catalytic Domain

Cat. No. MMP3-163H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human matrix metalloproteinase-3 (MMP-3, stromelysin-1, transin) cloned from human cDNA was expressed in <i>E.coli</i> . It consists of the catalytic domain of human MMP-3 (residues 105-265 swissprot accession P08254), and has been inactivated by mutagenesis (E219A). MW=18kDa.
Species	Human
Source	E.coli
ProteinLength	105-265 a.a.
Description	Stromelysin-1 is a protein that in humans is encoded by the MMP3 gene. Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP"s are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation.
Purity	> 95% by SDS-PAGE. The enzyme was observed as a single band migrating at a molecular weight of < 20kDa.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Specific Activity	Not detected. Activity described as U=100pmol/min at 37°C using a colorimetric assay with thiopeptolide Ac-Pro-Leu-Gly-[2-mercapto-4-methyl-pentanoyl]-Leu-Gly-OC ₂ H ₅ (Biomol) as substrate.
Usage	Enzyme kinetic studies, cleavage of target substrates and screening of inhibitors.
Supplied As	0.2mg/ml in 20mM Tris, pH7.2, 10mM CaCl ₂ , 0.1mM ZnCl ₂ , 0.3M NaCl, 0.2M Acetohydroxamic Acid (AHA). The concentration is calculated from the absorbance at 280nm, (e ₂₈₀ =27310M ⁻¹ cm ⁻¹).
Note	Under the above described conditions, to avoid precipitation or protein dimerization, the product can be concentrated to a maximum of 0.5mM.
Storage	-80°C. The enzyme is stable at -20°C for at least 1 week. After initial defrost, aliquot enzyme into individual tubes and refreeze at -80°C. Avoid repeated freeze/thaw cycles.

GENE INFORMATION

Gene Name	MMP3 matrix metalloproteinase 3 (stromelysin 1, procollagenase) [Homo sapiens]
Synonyms	MMP3; matrix metalloproteinase 3 (stromelysin 1, procollagenase); matrix metalloproteinase 3 (stromelysin 1, procollagenase); SL-1; STMY; STR1; CHDS6; MMP-3; STMY1; MGC126102; MGC126103; MGC126104; matrix metalloproteinase 3; Matrix metalloproteinase-3; transin-1; Stromelysin-1; proteoglycanase; EC 3.4.24.17; Transin-1
Gene ID	4314
mRNA Refseq	NM_002422

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_002413
MIM	185250
UniProt ID	P08254
Chromosome Location	11q22.3
Function	calcium ion binding; metalloendopeptidase activity; peptidase activity; zinc ion binding
PDB rendering based on 1b3d.	