

Recombinant Mouse Prpf19 Protein, Myc/DDK-tagged

Cat. No. Prpf19-5145M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Purified recombinant protein of mouse full-length pre-mRNA processing factor 19 (Prpf19), with C-terminal MYC/DDK tag, expressed in HEK293T cells.
Species	Mouse
Source	HEK293
Description	Ubiquitin-protein ligase which is a core component of several complexes mainly involved in pre-mRNA splicing and DNA repair. Required for pre-mRNA splicing as component of the spliceosome. Core component of the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome and participates in its assembly, its remodeling and is required for its activity. During assembly of the spliceosome, mediates 'Lys-63'-linked polyubiquitination of the U4 spliceosomal protein PRPF3. Ubiquitination of PRPF3 allows its recognition by the U5 component PRPF8 and stabilizes the U4/U5/U6 tri-snRNP spliceosomal complex. Recruited to RNA polymerase II C-terminal domain (CTD) and the pre-mRNA, it may also couple the transcriptional and spliceosomal machineries. The XAB2 complex, which contains PRPF19, is also involved in pre-mRNA splicing, transcription and transcription-coupled repair. Beside its role in pre-mRNA splicing PRPF19, as part of the PRP19-CDC5L complex, plays a role in the DNA damage response/DDR. It is recruited to the sites of DNA damage by the RPA complex where PRPF19 directly ubiquitinates RPA1 and RPA2. 'Lys-63'-linked polyubiquitination of the RPA complex allows the recruitment of the ATR-ATRIP complex and the activation of ATR, a master regulator of the DNA damage response. May also play a role in DNA double-strand break

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(DSB) repair by recruiting the repair factor SETMAR to altered DNA. As part of the PSO4 complex may also be involved in the DNA interstrand cross-links/ICLs repair process. In addition, may also mediate 'Lys-48'-linked polyubiquitination of substrates and play a role in proteasomal degradation May play a role in the biogenesis of lipid droplets May play a role in neural differentiation possibly through its function as part of the spliceosome.

Molecular Mass	55.7 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Stability	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Storage	Store at -80 centigrade after receiving vials.
Concentration	>50 µg/mL as determined by microplate BCA method
Storage Buffer	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.

GENE INFORMATION

Gene Name	Prpf19 pre-mRNA processing factor 19 [<i>Mus musculus</i> (house mouse)]
Official Symbol	Prpf19
Synonyms	PRPF19; PRP19/PSO4 pre-mRNA processing factor 19 homolog (<i>S. cerevisiae</i>); pre-mRNA-processing factor 19; prp19 beta; PRP19/PSO4 homolog; senescence evasion factor; nuclear matrix protein 200; nuclear matrix protein SNEV; PSO4; Snev; Prp19; NMP200; AA617263; AL024362; D19Wsu55e

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	28000
mRNA Refseq	NM_134129
Protein Refseq	NP_598890
UniProt ID	Q99KP6

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA