

Active Recombinant Human CD46 Protein, Fc-tagged

Cat. No. CD46-3963H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD46 Protein(Cys35-Asp328) with a human IgG1 (Pro100-Lys330) Fc tag at C-terminus was expressed in Chinese Hamster Ovary cell line.
Species	Human
Source	CHO
ProteinLength	35-328
Description	The protein encoded by this gene is a type I membrane protein and is a regulatory part of the complement system. The encoded protein has cofactor activity for inactivation of complement components C3b and C4b by serum factor I, which protects the host cell from damage by complement. In addition, the encoded protein can act as a receptor for the Edmonston strain of measles virus, human herpesvirus-6, and type IV pili of pathogenic Neisseria. Finally, the protein encoded by this gene may be involved in the fusion of the spermatozoa with the oocyte during fertilization. Mutations at this locus have been associated with susceptibility to hemolytic uremic syndrome. Alternatively spliced transcript variants encoding different isoforms have been described.
Form	Disulfide-linked homodimer
Bio-activity	Measured by its binding ability in a functional ELISA. When Recombinant Human CD46 Fc Chimera is immobilized at 5 µg/mL (100 µL/well), the concentration of Human C4b that produces 50% of the optimal binding response is 2-12

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	µg/mL.
Molecular Mass	59 kDa
N-terminal Sequence Analysis	Cys35
Endotoxin	<0.10 EU/µg of the protein by the LAL method.
Purity	> 95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <p>12 months from date of receipt, -20 to -70 centigrade as supplied.</p> <p>1 month, 2 to 8 centigrade under sterile conditions after reconstitution.</p> <p>3 months, -20 to -70 centigrade under sterile conditions after reconstitution.</p>
Storage Buffer	Lyophilized from a 0.2 µm filtered solution in PBS.
Reconstitution	Reconstitute at 500 µg/mL in PBS.
Shipping	The product is shipped at ambient temperature.

GENE INFORMATION

Gene Name	CD46 CD46 molecule [Homo sapiens (human)]
Official Symbol	CD46
Synonyms	CD46; CD46 molecule; MCP; TLX; AHUS2; MIC10; TRA2.10; membrane cofactor protein; CD46 antigen, complement regulatory protein; CD46 molecule, complement

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

regulatory protein; antigen identified by monoclonal antibody TRA-2-10; complement membrane cofactor protein; measles virus receptor; membrane cofactor protein (CD46, trophoblast-lymphocyte cross-reactive antigen); trophoblast leucocyte common antigen; trophoblast leucocyte common antigen; trophoblast-lymphocyte cross-reactive antigen

Gene ID [4179](#)


mRNA Refseq [NM_172361](#)

Protein Refseq [NP_758871](#)

MIM [120920](#)

UniProt ID [P15529](#)

Binding Activity




When Recombinant Human CD46 Fc Chimera is immobilized at 5 µg/mL (100 µL/well), it binds to Human C4b with an ED50 of 2-12 µg/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



2 µg/lane of Recombinant Human CD46 Fc Chimera was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 88-98 kDa and 160-190 kDa, respectively.