

Recombinant Human ENO1 Protein

Cat. No. ENO1-022H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant full length human ENO1 protein (1-434) without tag was purified from E. coli citrullinated by PAD4.
Species	Human
Source	E.coli
ProteinLength	1-434
Description	This gene encodes alpha-enolase, one of three enolase isoenzymes found in mammals. Each isoenzyme is a homodimer composed of 2 alpha, 2 gamma, or 2 beta subunits, and functions as a glycolytic enzyme. Alpha-enolase in addition, functions as a structural lens protein (tau-crystallin) in the monomeric form. Alternative splicing of this gene results in a shorter isoform that has been shown to bind to the c-myc promoter and function as a tumor suppressor. Several pseudogenes have been identified, including one on the long arm of chromosome 1. Alpha-enolase has also been identified as an autoantigen in Hashimoto encephalopathy.
Molecular Mass	47.74 kDa
Purity	≥90% as determined by SDS-PAGE
Stability	≥ 6 months

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage At -80 centigrade.

Storage Buffer TBS, pH 7.4

GENE INFORMATION

Gene Name ENO1 enolase 1 [Homo sapiens (human)]

Official Symbol ENO1

Synonyms ENO1; enolase 1; NNE; PPH; MPB1; ENO1L1; HEL-S-17; alpha-enolase; c-myc promoter-binding protein-1; 2-phospho-D-glycerate hydro-lyase; MYC promoter-binding protein 1; alpha enolase like 1; enolase 1, (alpha); enolase-alpha; epididymis secretory protein Li 17; non-neural enolase; phosphopyruvate hydratase; plasminogen-binding protein; tau-crystallin; EC 4.2.1.11

Gene ID 2023

mRNA Refseq NM_001428

Protein Refseq NP_001419

MIM 172430


UniProt ID P06733

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**SDS-PAGE analysis
of citrullinated α -
enolase**




Lane 1: MW Markers

Lane 2: 4 μ g α -enolase (PAD4 Citrullinated)

Lane 3: 2 μ g α -enolase (PAD4 Citrullinated)

**WB analysis of α -
enolase citrullination**



Lane 1: Citrullinated α -enolase (20 ng)

Lane 2: Citrullinated α -enolase (100 ng)

Lane 3: Citrullinated α -enolase (400 ng)

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Lane 4: α -enolase (20 ng)

Lane 5: α -enolase (100 ng)

Lane 6: α -enolase (400 ng)

α -enolase and Citrullinated α -enolase were reacted with Citrullinated α -enolase Monoclonal Antibody and detected using Goat Anti-Mouse IgG HRP.

Identification of modified sites in Citrullinated α -enolase.

MSILKIHARE	IFDSRGNPTV	EVDLFTSKGL	FRAAVPSGAS	TGIYEALER
DNDKTRYMGK	GVSKAVEHIN	KTIAPALVSK	KLVNTEQEKI	DKLMIEMDGT
ENKSKFGANA	ILGVSLAVCK	AGAVEKGVPL	YRHIADLAGN	SEVILPVPAF
NVINGGSHAG	NKLAMQEFMI	LPVGAANFRE	AMRIGAEVYH	NLKNVIKEY
GKDATNVGDE	GGFAPNILEN	KEGLELLKTA	IGKAGYTDKV	VIGMDVAASE
FFRSGKYDLD	FKSPDDPSRY	ISPDQLADLY	KSFIKDYPPV	SIEDPFDQDD
WGAWQKFTAS	AGIQVVGDDL	TVTNPKRIAK	AVNEKSCNCL	LLKVNQIGSV
TESLQACKLA	QANGWGMVMS	HRSGETEDTF	IADLVVGLCT	GQIKTGAPCR
SERLAKYNQL	LRIEEELGSK	AKFAGRNFN	PLAK	

Citrullinated α -enolase was detected by LC-MS/MS and analyzed using Mascot and Scaffold PTM sofware. Deiminated arginines are indicated in teal.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA