

Recombinant Human PCSK9 Protein, His-tagged

Cat. No. PCSK9-135H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active recombinant N-terminal His-tagged human PCSK9 protein (31-692) isolated from a HEK293 overexpression system.
Species	Human
Source	HEK293
ProteinLength	31-692
Description	This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an autocatalytic processing event with its prosegment in the ER and is constitutively secreted as an inactive protease into the extracellular matrix and trans-Golgi network. It is expressed in liver, intestine and kidney tissues and escorts specific receptors for lysosomal degradation. It plays a role in cholesterol and fatty acid metabolism. Mutations in this gene have been associated with autosomal dominant familial hypercholesterolemia. Alternative splicing results in multiple transcript variants.
Form	Lyophilized
Molecular Mass	71 kDa (~16 & 63 kDa purified)
Purity	≥90% as determined by SDS-PAGE

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Stability	≥ 6 months
Storage	At -80 centigrade.
Storage Buffer	PBS, pH 7.4, with 30% sucrose

GENE INFORMATION


Gene Name	PCSK9 proprotein convertase subtilisin/kexin type 9 [Homo sapiens (human)]
Official Symbol	PCSK9
Synonyms	PCSK9; proprotein convertase subtilisin/kexin type 9; FH3; PC9; FHCL3; NARC1; LDLCQ1; NARC-1; HCHOLA3; proprotein convertase subtilisin/kexin type 9; convertase subtilisin/kexin type 9 preproprotein; neural apoptosis regulated convertase 1; subtilisin/kexin-like protease PC9
Gene ID	255738
mRNA Refseq	NM_174936
Protein Refseq	NP_777596
MIM	607786
UniProt ID	Q8NBP7

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE analysis of PCSK9



Lane 1: MW Ladder

Lane 2: PCSK9 (1 µg)

Lane 3: PCSK9 (2 µg)

PCSK9 inhibits LDL uptake.

