Active Recombinant Human FLT4 protein, Tyr25-Glu775-tagged, Biotinylated

Cat. No. FLT4-051H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human FLT4(Tyr25-Glu775) protein, fused to His/Avi tag at the C-terminus, was expressed in HEK293 cells .
Species	Human
Source	HEK293
ProteinLength	Tyr25-Glu775
Description	VEGFR2 (KDR/Flk-1), VEGFR1 (Flt-1) and VEGFR3 (Flt-4) belong to the class III subfamily of receptor tyrosine kinases (RTKs). All three receptors contain seven immunoglobulin-like repeats in their extracellular domains and kinase insert domains in their intracellular regions. The expression of VEGFR1, 2, and 3 is almost exclusively restricted to the endothelial cells. These receptors are likely to play essential roles in vasculogenesis and angiogenesis. VEGFR3 cDNA encodes a 1298 amino acid (aa) residue precursor protein with a 24 aa residue signal peptide. Mature VEGFR3 is composed of a 751 aa residue extracellular domain, a 22 aa residue transmembrane domain and a 482 aa residue cytoplasmic domain. Both VEGF-C and VEGF-D have been shown to bind and activate VEGFR3 (Flt-4). VEGFR3 is widely expressed in the early embryo but becomes restricted to lymphatic endothelia at later stages of development. It is likely that VEGFR3 may be important for lymph angiogenesis.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Predicted N Terminal	Tyr25
Form	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.
Bio-activity	The biotin to protein ratio is greater than 0.7 as determined by the HABA assay. Measured by its binding ability in a functional ELISA. When Recombinant Human VEGF-C is immobilized at 0.5 g/mL (100 μ/well), Biotinylated Recombinant Human FLT4 His-tag Avi-tag binds with an ED50 of 15-90 ng/mL.
Molecular Mass	115-140 kDa, 72-89 kDa and 60-70 kDa
Endotoxin	<0.10 EU per 1 μg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Applications	Bioactivity
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.12 months from date of receipt, -20 to -70 °C as supplied.1 month, 2 to 8 °C under sterile conditions after reconstitution.3 months, -20 to -70 °C under sterile conditions after reconstitution.
Reconstitution	Reconstitute at 500 μg/mL in PBS.
Conjugation	Biotin

GENE INFORMATION

Gene Name	FLT4 fms-related tyrosine kinase 4 [Homo sapiens]

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	FLT4
Synonyms	FLT4; fms-related tyrosine kinase 4; vascular endothelial growth factor receptor 3; PCL; VEGFR3; FLT-4; VEGFR-3; soluble VEGFR3 variant 1; soluble VEGFR3 variant 2; soluble VEGFR3 variant 3; fms-like tyrosine kinase 4; tyrosine-protein kinase receptor FLT4; FLT41; LMPH1A;
Gene ID	2324
mRNA Refseq	NM_002020
Protein Refseq	NP_002011
MIM	136352
UniProt ID	P35916
Binding Activity	

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA