

Active Recombinant Human HAVCR2 protein, His/Avi-tagged, Biotinylated

Cat. No. HAVCR2-051H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human HAVCR2(Ser22-Arg200) protein, fused to His/Avi tag at the C-terminus, was expressed in CHO cells .
Species	Human
Source	CHO
ProteinLength	Ser22-Arg200
Description	HAVCR2 is a 60 kDa member of the TIM family of immune regulating molecules. TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk region (1, 2). Mature human TIM-3 consists of a 181 amino acid (aa) extracellular domain (ECD), a 21 aa transmembrane segment, and a 78 aa cytoplasmic tail (3). An alternatively spliced isoform is truncated within the mucin-like stalk. Within the ECD, human TIM-3 shares 58% aa sequence identity with mouse and rat TIM-3. TIM-3 is up-regulated on several populations of activated myeloid cells (macrophage, monocyte, dendritic cell, microglia, mast cell) and T cells (Th1, CD8+, NK, Treg) (3-10). Its binding to Galectin-9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor immunity (11). TIM-3 ligation attenuates CD8+ and Th1 cell responses (11-13) and promotes the activity of Treg and myeloid derived suppressor cells (8, 11, 13, 14). In addition, dendritic cell-expressed TIM-3 dampens inflammation by enabling the phagocytosis of apoptotic cells and the cross-presentation of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

apoptotic cell antigens (4). It also binds the alarmin HMGB1, thereby preventing the activation of TLRs in response to released tumor cell DNA (7). TIM-3 interactions with Galectin-9 can alternatively trigger immune stimulatory effects, such as the coactivation of NK cell cytotoxicity (10).

Predicted N Terminal	Ser22
Form	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
Bio-activity	Measured by its binding ability in a functional ELISA. When Recombinant Human Galectin9 is immobilized at 1 µg/mL (100 µL/well), the concentration that produces 50% of the optimal binding response is 0.1-0.8 µg/mL.
Molecular Mass	37-50 kDa, under reducing conditions
Endotoxin	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Applications	Bioactivity
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Reconstitution	Reconstitute at 200 µg/mL in PBS.
Conjugation	Biotin

GENE INFORMATION

Gene Name HAVCR2 hepatitis A virus cellular receptor 2 [Homo sapiens]

Official Symbol HAVCR2

Synonyms HAVCR2; hepatitis A virus cellular receptor 2; FLJ14428; Tim 3; TIM3; TIMD3; kidney injury molecule-3; T-cell membrane protein 3; T cell immunoglobulin mucin 3; T cell immunoglobulin mucin-3; T-cell immunoglobulin and mucin domain-containing protein 3; KIM-3; Tim-3; TIMD-3; HAVcr-2;

Gene ID 84868


mRNA Refseq NM_032782

Protein Refseq NP_116171

MIM 606652

UniProt ID Q8TDQ0

Binding Activity




 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA