Recombinant Human IL8, His-tagged

Cat. No. IL8-11727H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL8 protein, fused to His-tag, was expressed in E.coli and purified by Ni-sepharose.
Species	Human
Source	E.coli
ProteinLength	1-99a.a.
Description	The protein encoded by this gene is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. This gene and other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region mapped to chromosome 4q.
Storage	The protein is stored in PBS buffer at -20℃. Avoid repeated freezing and thawing cycles.
Storage Buffer	1M PBS (58mM Na2HPO4,17mM NaH2PO4, 68mM NaCl, pH8.) added with 300mM Imidazole and 0.7% Sarcosyl, 15%glycerol.

GENE INFORMATION

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	IL8 interleukin 8 [Homo sapiens]
Official Symbol	IL8
Synonyms	IL8; interleukin 8; interleukin-8; 3 10C; alveolar macrophage chemotactic factor I; AMCF I; b ENAP; beta endothelial cell derived neutrophil activating peptide; chemokine (C X C motif) ligand 8; CXCL8; GCP 1; GCP1; granulocyte chemotactic protein 1; IL 8; K60; LECT; LUCT; lung giant cell carcinoma derived chemotactic protein; lymphocyte derived neutrophil activating peptide; LYNAP; MDNCF; MONAP; monocyte derived neutrophil chemotactic factor; monocyte derived neutrophil activating peptide; NAF; NAP 1; NAP1; neutrophil activating peptide 1; SCYB8; TSG 1; tumor necrosis factor induced gene 1; emoctakin; T-cell chemotactic factor; neutrophil-activating peptide 1; chemokine (C-X-C motif) ligand 8; beta-thromboglobulin-like protein; tumor necrosis factor-induced gene 1; monocyte-derived neutrophil chemotactic factor; monocyte-derived neutrophil-activating peptide; small inducible cytokine subfamily B, member 8; lung giant cell carcinoma-derived chemotactic protein; beta endothelial cell-derived neutrophil activating peptide; GCP-1; NAP-1;
Gene ID	3576
mRNA Refseq	NM_000584
Protein Refseq	NP_000575
MIM	146930
UniProt ID	P10145
Chromosome Location	4q13-q21

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	ATF-2 transcription factor network, organism-specific biosystem; Activation of Genes by ATF4, organism-specific biosystem; Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem;
Function	chemokine activity; interleukin-8 receptor binding; protein binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA