

Active Recombinant Bovine Interleukin 23, Alpha Subunit P19

Cat. No. IL23A-239B **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Bovine IL-23 alpha subunit p19 was expressed in Yeast. MW= 19.0 kDa (171 aa).
Species	Bovine
Source	Yeast
Description	Interleukin-23 subunit alpha is a protein that in humans is encoded by the IL23A gene. This gene encodes the p19 subunit of the heterodimeric cytokine interleukin 23 (IL23). IL23 is composed of this protein and the p40 subunit of interleukin 12 (IL12B). The receptor of IL23 is formed by the beta 1 subunit of IL12 (IL12RB1) and an IL23 specific subunit, IL23R. Both IL23 and IL12 can activate the transcription activator STAT4, and stimulate the production of interferon-gamma (IFNG). In contrast to IL12, which acts mainly on naive CD4(+) T cells, IL23 preferentially acts on memory CD4(+) T cells.
Formulation/ Reconstitution	Lyophilized. Reconstitute with sterile phosphate-buffered saline containing at least 0.1% carrier protein.
Purification	Ion-exchange chromatography.
Purity	>95% as visualized by SDS-PAGE analysis.
Amino Acid Sequence	RAVSEDSSPA WTRGQQLSQQ LCMLAWSAHL PMGHMDLPRE EGGDKTTDDV PRIQCEDGCD PQGLRDNSQS CLQRIHRLV FYEKLIGSDI FTGEPSSLPN

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GPVDQLHASI LGLRELLQPK GHHWETEQTP SPIPSQPWQR LLLRLKILRS
LQAFVAVAAAR VFAHGAATLS P.

Bioactivity

The biological activity of recombinant bovine IL-23, alpha subunit p19 has not been tested. The active form of IL-23 is a dimer (p19/p40).

Stability And Storage

Stable for up to twelve months from date of receipt at -20°C. Stable for at least 3 months when stored in working aliquots with a carrier protein at -20°C. Avoid repeated freeze/thaw cycles.

GENE INFORMATION

Gene Name

IL23A interleukin 23, alpha subunit p19 [Bos taurus]

Synonyms

interleukin 23, alpha subunit p19; IL23A; IL-23

Gene ID

511022

mRNA Refseq

[XM_588269.4](#)

Protein Refseq

[XP_588269.2](#)

**Chromosome
Location**

5

Pathway

Cytokine-cytokine receptor interaction; Jak-STAT signaling pathway

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA