
Active Recombinant Chicken TIMP2 protein, His-tagged

Cat. No. TIMP2-411C **Lot. No.** (See product label)


SPECIFICATION

Product Overview	Active Recombinant Chicken TIMP2 protein(O42146)(27~220 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Chicken
Source	E.coli
ProteinLength	27-220 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Tissue inhibitors of metalloproteinases or TIMPs are a family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs). There are four members of the family, TIMP-1, TIMP-2, TIMP-3, and TIMP-4. TIMP-2 is a non N-glycosylated protein with a molecular mass of 22 kDa produced by a wide range of cell types, which inhibits MMPs non-covalently by the formation of binary complexes. TIMP-2 also has erythroid potentiating and cell growth promoting activities. The activity of recombinant chicken TIMP2 was measured by its ability to inhibit rhMMP2 cleavage of a fluorogenic peptide substrate MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH2 in the assay buffer 50 mM Tris, 10 mM CaCl2, 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. rhMMP2 was diluted to 100 ug/ml and activated with 1 mM APMA at 37°C for 1 hour and rgTIMP2 (MW: 23 KD) was diluted to different concentrations with the assay buffer. Mix 8 µof rgTIMP2 curve dilutions, 12.8 µof activated rhMM

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	P-2, and 59.2 µof assay buffer, including a control containing assay buffer and the dilu ted rhMMP-2 and incubate the reactions for 2 hours at 37°C. Loading 50 µof the incub ated mixtures which were diluted five-fold in assay buffer into empty wells of a plate, a nd start the reaction by adding 50 µof 20 µM substrate. Include a substrate blank contai ning 50 µof assay buffer and 50 µof 20 µM substrate. Then read at excitiation and emis sion wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The result was shown and it was obvious that recombinant chicken TIMP2 significantl y decreased rhMMP2 activity. The inhibition IC50 was <16 nM.
Molecular Mass	25.4kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
Activity	

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE	

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA