

Active Recombinant Human CD40 Protein, Fc-tagged, Alexa Fluor 555 conjugated

Cat. No. CD40-108HAF555 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Alexa Fluor 555 conjugated recombinant human CD40 Protein extracellular domain 21-193aa (signal peptide removed), fused with a C-terminal Fc tag, was expressed in CHO.
Species	Human
Source	CHO
ProteinLength	405
Description	CD40, also known as TNFRSF5, is a member of the TNF receptor superfamily which are single transmembrane-spanning glycoproteins. CD40 protein plays an essential role in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. CD40 protein is expressed in B cells, dendritic cells, macrophages, endothelial cells, and several tumor cell lines. Defects in CD40 result in hyper-IgM immunodeficiency type 3 (HIGM3). In addition, CD40/CD40L interaction is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis.
Form	Lyophilized
Bio-activity	Measured by its binding ability in a functional ELISA.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	45.2 kDa
N-terminal Sequence Analysis	Glu 21
Endotoxin	< 1.0 EU/ µg by the LAL method.
Purity	> 95 % as determined by SDS-PAGE
Characteristic	Disulfide-linked homodimer Labeled with Alexa Fluor 555 via amines With an excitation and emission maximum of 555/565 nm, Alexa Fluor 555 can be efficiently excited using a 543 nm He-Ne laser line and detected under standard TRITC/Cy3 filters.
Stability	Samples are stable for up to 12 months from date of receipt at -70 centigrade.
Storage	Store it under sterile conditions at -20 to -70 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Lyophilized from sterile PBS, pH 7.4.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.
Conjugation	Alexa Fluor 555

GENE INFORMATION

Gene Name	CD40 CD40 molecule, TNF receptor superfamily member 5 [Homo sapiens]
------------------	--

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	CD40
Synonyms	CD40; CD40 molecule, TNF receptor superfamily member 5; TNFRSF5, tumor necrosis factor receptor superfamily, member 5; tumor necrosis factor receptor superfamily member 5; Bp50; p50; CD40L receptor; CD40 type II isoform; B cell-associated molecule; B cell surface antigen CD40; B-cell surface antigen CD40; CD40 antigen (TNF receptor superfamily member 5); tumor necrosis factor receptor superfamily, member 5; nerve growth factor receptor-related B-lymphocyte activation molecule; CDW40; TNFRSF5; MGC9013;
Gene ID	958
mRNA Refseq	NM_001250
Protein Refseq	NP_001241
MIM	109535
UniProt ID	P25942

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA