

Recombinant Monkey CD80 Protein, hFc-tagged, Alexa Fluor 555 conjugated

Cat. No. CD80-137CAF555 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Alexa Fluor 555 conjugated recombinant Cynomolgus/Rhesus macaque CD80 (Val35-Asn242) protein was fused to human IgG1 Fc tag at C-terminus and expressed in human 293 cells (HEK293).
Species	Monkey
Source	HEK293
ProteinLength	Val35-Asn242
Description	B7-1 and B7-2, together with their receptors CD28 and CTLA4, constitute one of the dominant co-stimulatory pathways that regulate T and B cell responses. Although both CTLA4 and CD28 can bind to the same ligands, CTLA4 binds to B7-1 and B7-2 with a 20-100 fold higher affinity than CD28 and is involved in the downregulation of the immune response. B-lymphocyte activation antigen B7-1 (referred to as B7) also known as cluster of Differentiation 80 (CD80), is a member of cell surface immunoglobulin superfamily and is expressed on activated B cells, activated T cells, macrophages and dendritic cells. It is the ligand for two different proteins on the T cell surface: CD28 (for autoregulation and intercellular association) and CTLA-4 (for attenuation of regulation and cellular disassociation). CD80 works in tandem with CD86 to prime T cells. CD80 plays a role in induction of innate immune responses by activating NF-κB-signaling pathway in macrophages. CD80 is thus regarded as promising therapeutic targets for

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	autoimmune diseases and various carcinomas.
Form	Lyophilized
Molecular Mass	The protein has a calculated MW of 50.5 kDa. The protein migrates as 60-90 kDa under reducing (R) condition (SDS-PAGE) due to different glycosylation.
N-terminal Sequence Analysis	Val 35
Endotoxin	< 1.0 EU/ µg by the LAL method.
Purity	> 95 % as determined by SDS-PAGE
Characteristic	Disulfide-linked homodimer Labeled with Alexa Fluor 555 via amines With an excitation and emission maximum of 555/565 nm, Alexa Fluor 555 can be efficiently excited using a 543 nm He-Ne laser line and detected under standard TRITC/Cy3 filters.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.

Conjugation Alexa Fluor 555

GENE INFORMATION

Gene Name CD80

Official Symbol CD80

Synonyms CD80; CD80 antigen; T-lymphocyte activation antigen CD80; B7 protein; activation B7-1 antigen; B71; Ly53; TSA1; Cd28l; Ly-53; MIC17

Gene ID 102140078

mRNA Refseq XM_015446388

Protein Refseq XP_015301874

UniProt ID F7HRG1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA