

Recombinant Rat Ctla4 Protein, hFc-tagged, Alexa Fluor 647 conjugated

Cat. No. Ctla4-1143RAF647 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Alexa Fluor 647 conjugated recombinant Rat Ctla4 (Ile38-Asp161) protein was fused to human IgG1 Fc tag at C-terminus and expressed in human 293 cells (HEK293).
Species	Rat
Source	HEK293
ProteinLength	Ile38-Asp161
Description	CTLA-4 (Cytotoxic T-Lymphocyte Antigen 4) is also known as CD152 (Cluster of differentiation 152), is a protein receptor that downregulates the immune system. CTLA4 is a member of the immunoglobulin superfamily, which is expressed on the surface of Helper T cells and transmits an inhibitory signal to T cells. The protein contains an extracellular V domain, a transmembrane domain, and a cytoplasmic tail. Alternate splice variants, encoding different isoforms. CTLA4 is similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to CD80 and CD86, also called B7-1 and B7-2 respectively, on antigen-presenting cells. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may be important to their function. Fusion proteins of CTLA4 and antibodies (CTLA4-Ig) have been used in clinical trials for rheumatoid arthritis.
Form	Lyophilized

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	The protein has a calculated MW of 40.2 kDa. The protein migrates as 50-55 kDa under reducing (R) condition (SDS-PAGE).
N-terminal Sequence Analysis	Ile 38
Endotoxin	< 1.0 EU/ µg by the LAL method.
Purity	> 95 % as determined by SDS-PAGE
Characteristic	Disulfide-linked homodimer Labeled with Alexa Fluor 647 via amines Excitation = 650 nm Emission = 668 nm
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.
Conjugation	Alexa Fluor 647

GENE INFORMATION

Gene Name	Ctla4
Official Symbol	Ctla4
Synonyms	CTLA4; cytotoxic T-lymphocyte-associated protein 4; cytotoxic T-lymphocyte protein 4; soluble form; CD152 antigen; transmembrane form; Cd152; CTLA-4; sCTLA4
Gene ID	63835
mRNA Refseq	NM_031674
Protein Refseq	NP_113862
UniProt ID	Q62859

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA