

Recombinant Human CD79B Protein, hFc-tagged, FITC conjugated

Cat. No. CD79B-239HF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	FITC conjugated recombinant human CD79B protein (Ala29-Asp159), fused to human IgG1 Fc tag at C-terminus, was expressed in human 293 cells (HEK293).
Species	Human
Source	HEK293
ProteinLength	131
Description	B-cell antigen receptor complex-associated protein beta chain (CD79b) is also known as B-cell-specific glycoprotein B29, Ig-beta, Immunoglobulin-associated B29 protein, B 29 and IGB, which is a single-pass type I membrane protein containing one Ig-like V-type (immunoglobulin-like) domain and one ITAM domain. CD79b is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR). CD79b can enhance phosphorylation of CD79A, possibly by recruiting kinases which phosphorylate CD79A or by recruiting proteins which bind to CD79A and protect it from dephosphorylation. Defects in CD79b are the cause of agammaglobulinemia type 6 (AGM6) that is a primary immunodeficiency characterized by profoundly low or absent serum antibodies and low or absent circulating B cells due to an early block of B-cell development.
Form	Lyophilized
Molecular Mass	The protein has a calculated MW of 41.8 kDa. The protein migrates as 50-65 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	< 1.0 EU/ µg by the LAL method.
Purity	> 92 % as determined by SDS-PAGE
Characteristic	Disulfide-linked homodimer Labeled with FITC via amines Excitation source: 488 nm spectral line, argon-ion laser Excitation Wavelength: 488 nm Emission Wavelength: 535 nm
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.
Conjugation	FITC

GENE INFORMATION

Gene Name **CD79B**

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	CD79B
Synonyms	CD79B; CD79b molecule, immunoglobulin-associated beta; CD79B antigen (immunoglobulin associated beta) , IGB; B-cell antigen receptor complex-associated protein beta chain; B29; Ig-beta; B-cell-specific glycoprotein B29; immunoglobulin-associated B29 protein; CD79b antigen (immunoglobulin-associated beta); IGB; AGM6
Gene ID	974
mRNA Refseq	NM_000626
Protein Refseq	NP_000617
MIM	147245
UniProt ID	P40259

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA