

Active Recombinant Monkey MET Protein, Fc-tagged, FITC conjugated

Cat. No. MET-183CF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	A DNA sequence encoding the Monkey /rhesus MET (NP_001162100.1) (Met1-Thr932) (FITC conjugated) was expressed with the Fc region of human IgG1 at the C-terminus. Monkey and Rhesus MET sequences are identical.
Species	Monkey
Source	HEK293
ProteinLength	Met1-Thr932 1146
Form	Lyophilized
Bio-activity	Immobilized Cynomolgus HGF at 10 µg/mL (100 µL/well) can bind Cynomolgus MET-Fc, EC50 of Cynomolgus MET-Fc is 0.04-0.09 µg/mL.
Molecular Mass	The recombinant cynomolgus/rhesus MET comprises 1146 amino acids and has a calculated molecular mass of 128.3 kDa. The apparent molecular mass of it is approximately 100.2 and 42.5 kDa in SDS-PAGE under reducing conditions.
Endotoxin	< 1.0 EU/ µg of the protein as determined by the LAL method.
Purity	(66.7+31.5) % as determined by SDS-PAGE
Characteristic	Disulfide-linked homodimer

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	Labeled with FITC via amines Excitation source: 488 nm spectral line, argon-ion laser Excitation Wavelength: 488 nm Emission Wavelength: 535 nm
Stability	Samples are stable for up to 12 months from date of receipt at -70 centigrade.
Storage	Store it under sterile conditions at -70 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Lyophilized from sterile PBS, pH7.4.
Conjugation	FITC
GENE INFORMATION	
Gene Name	METmet proto-oncogene (hepatocyte growth factor receptor) [Macaca mulatta(Rhesus monkey)]
Official Symbol	MET
Synonyms	MET; met proto-oncogene (hepatocyte growth factor receptor); hepatocyte growth factor receptor; NP_001162100.1; EC 2.7.10.1
Gene ID	704562
mRNA Refseq	NM_001168629
Protein Refseq	NP_001162100