

Recombinant Human TNFRSF17 Protein, Fc/His-tagged, FITC conjugated

Cat. No. TNFRSF17-2348HF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	FITC conjugated recombinant human TNFRSF17 fused with Fc/His tag at C-terminal was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	302
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation.
Form	Lyophilized
Molecular Mass	The recombinant human TNFRSF17/Fc is a disulfide-linked homodimer. The reduced monomer comprises 302 amino acids after removal of the signal peptide and has a predicted molecular mass of 34 kDa. As a result of glycosylation, the apparent molecular mass of rh TNFRSF17/Fc monomer is approximately 40 kDa in SDS-PAGE under reducing conditions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	< 1.0 EU/ µg of the protein as determined by the LAL method.
Characteristic	Disulfide-linked homodimer Labeled with FITC via amines Excitation source: 488 nm spectral line, argon-ion laser Excitation Wavelength: 488 nm Emission Wavelength: 535 nm
Stability	Samples are stable for up to 12 months from date of receipt at -70 centigrade.
Storage	Store it under sterile conditions at -20 to -70 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Storage Buffer	Lyophilized from sterile PBS, pH 7.4
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.
Conjugation	FITC

GENE INFORMATION

Gene Name	TNFRSF17 tumor necrosis factor receptor superfamily, member 17 [Homo sapiens]
Official Symbol	TNFRSF17
Synonyms	TNFRSF17; tumor necrosis factor receptor superfamily, member 17; BCMA; tumor necrosis factor receptor superfamily member 17; BCM; CD269; B-cell maturation factor; B cell maturation antigen; B-cell maturation protein;
Gene ID	608

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_001192
Protein Refseq	NP_001183
MIM	109545
UniProt ID	Q02223

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA