

Active Recombinant Human TNFRSF8 Protein, Fc-tagged, FITC conjugated

Cat. No. TNFRSF8-1594HF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	FITC conjugated recombinant human CD30 (amino acid residues 1 – 379) encoding the extracellular domain was fused to the carboxy-terminal polyhistidine tagged Fc region of human IgG1 via a polypeptide linker. The chimeric protein was expressed in <i>NSO cells</i> .
Species	Human
Source	Mammalian Cells
ProteinLength	1-379 a.a.
Description	CD30/TNFRSF8, a type I transmembrane glycoprotein belonging to the TNF receptor superfamily and tumor marker. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. It is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. CD30 is associated with anaplastic large cell lymphoma. It is expressed in embryonal carcinoma but not in seminoma and is thus is a useful marker in distinguishing between these germ cell tumors.
Form	Lyophilized

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Bio-activity	The biological activity of Human CD30 was determined by its ability to bind rhCD30 Ligand. Immobilized rhCD30 Ligand at 100 ng/mL (100 µL/well) can bind rhCD30/Fc with a linear range of 3.125-200 ng/mL in a functional ELISA.
Molecular Mass	100-125 kDa
Endotoxin	< 1.0 EU/ µg as determined by the LAL method
Purity	> 95 % by SDS-PAGE and analyzed by silver stain.
Characteristic	Disulfide-linked homodimer Labeled with FITC via amines Excitation source: 488 nm spectral line, argon-ion laser Excitation Wavelength: 488 nm Emission Wavelength: 535 nm
Stability	This lyophilized protein is stable for 6 to 12 months when stored desiccated at -20 to -70 centigrade. After aseptic reconstitution, this protein may be stored at 2 to 8 centigrade for one month or at -20 to -70 centigrade in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.
Storage	This lyophilized protein is stable for six to twelve months when stored desiccated at -20 to -70 centigrade. After aseptic reconstitution, this protein may be stored at 2 to 8 centigrade for one month or at -20 to -70 centigrade in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.
Storage Buffer	This recombinant protein was 0.2 µm filtered and lyophilized from modified Dulbecco's phosphate buffered saline (1X PBS) pH 7.2-7.3 with no calcium, magnesium, or preservatives.

Conjugation FITC

GENE INFORMATION

Gene Name TNFRSF8 tumor necrosis factor receptor superfamily, member 8 [Homo sapiens]

Official Symbol TNFRSF8

Synonyms

TNFRSF8; tumor necrosis factor receptor superfamily, member 8; CD30; Ki-1; D1S166E; Tumor necrosis factor receptor superfamily member 8; CD30L receptor; Lymphocyte activation antigen CD30; Ki-1 antigen; OTTHUMP00000001783; CD30 antigen

Gene ID 943

mRNA Refseq NM_001243

Protein Refseq NP_001234

MIM 153243

UniProt ID P28908

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA