

Active Recombinant Human IL3 protein, His-tagged

Cat. No. IL3-1036H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human IL3 protein(P08700)(20~152 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	20-152 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Interleukin 3 is an interleukin, a type of biological signal that can improve the body's natural response to disease as part of the immune system. It acts by binding to the interleukin 3 receptor IL-3 synergizes with other cytokines to stimulate the growth of immature progenitor cells of all lineages, and is therefore a multi-lineage colony-stimulating factor (CSF). It prevents cell death and promotes the survival of macrophages, mast cells, and megakaryocytes. To test the effect of IL-3 on cell proliferation, TF-1 cells were seeded into triplicate wells of 96-well plates at a density of 8,000 cells/well with 2% serum standard 1640 which contains various concentrations of recombinant human IL-3. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 μ l of CCK-8 solution was added to each well of the plate, then the absorbance at 450 nm was measured using a microplate reader after

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

incubating the plate for 2-4 hours at 37°C. Proliferation of TF-1 cells after incubation with IL-3 for 72h observed by inverted microscope was shown. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant human IL-3 for 72h. It was obvious that IL-3 significantly increased cell viability of TF-1 cells. The ED50 is 0.033-0.3 ng/ml.

Molecular Mass	16.7kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	IL3 interleukin 3 (colony-stimulating factor, multiple) [Homo sapiens]
Official Symbol	IL3
Synonyms	IL3; interleukin 3 (colony-stimulating factor, multiple); interleukin-3; hematopoietic growth factor; IL 3; mast cell growth factor; MCGF; MGC79398; MGC79399; MULTI CSF; multilineage colony stimulating factor; P cell stimulating factor; mast-cell growth factor; P-cell stimulating factor; P-cell-stimulating factor; multilineage-colony-stimulating factor; multipotential colony-stimulating factor; IL-3; MULTI-CSF;
Gene ID	3562

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq [NM_000588](#)

Protein Refseq [NP_000579](#)


MIM [147740](#)

UniProt ID [P08700](#)

Activity



SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA