

Active Recombinant Human IL18 protein, His-tagged

Cat. No. IL18-2296H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human IL18 protein(Q14116)(37~193 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	37-193 aa
Form	Lyophilized from sterile 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	Interleukin-18 (IL18, also known as interferon-gamma inducing factor) is a proinflammatory cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL18 works by binding to the interleukin-18 receptor, and together with IL-12 it induces cell-mediated immunity following infection with microbial products like lipopolysaccharide (LPS). Besides, Caspase 1 (CASP1) has been identified as an interactor of IL18, thus a binding ELISA assay was conducted to detect the interaction of recombinant human IL18 and recombinant human CASP1. Briefly, IL18 were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to CASP1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IL18 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50μ stop solution to the wells and read at 450nm immediately. The binding activity of IL18 and CASP1 was shown, and this effect was in a dose dependent manner.

Molecular Mass	19.5kDa
Endotoxin	<1.0EU per 1μg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (PH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	IL18 interleukin 18 (interferon-gamma-inducing factor) [Homo sapiens]
Official Symbol	IL18
Synonyms	IL18; interleukin 18 (interferon-gamma-inducing factor); interleukin-18; IGIF; IL 1g; IL 18; IL1F4; IL-1 gamma; iboctadekin; interleukin-1 gamma; IFN-gamma-inducing factor; IL-18; IL-1g; MGC12320;
Gene ID	3606
mRNA Refseq	NM_001243211
Protein Refseq	NP_001230140

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


MIM 600953

UniProt ID Q14116

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA