

Active Recombinant Human PDGFA protein, His-tagged

Cat. No. PDGFA-2570H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human PDGFA protein(P04085)(87~211 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	87-211 aa
Form	Lyophilized from sterile 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	Platelet Derived Growth Factor Subunit A (PDGFA) is a member of Platelet-derived growth factor (PDGF) family which plays a significant role in blood vessel formation, the growth of blood vessels from already-existing blood vessel tissue, mitogenesis, i.e. proliferation, of mesenchymal cells such as fibroblasts, osteoblasts, tenocytes, vascular smooth muscle cells and mesenchymal stem cells as well as chemotaxis, the directed migration, of mesenchymal cells. Besides, Platelet Derived Growth Factor Receptor Alpha (PDGFRa) has been identified as an interactor of PDGFA, thus a binding ELISA assay was conducted to detect the interaction of recombinant human PDGFA and recombinant human PDGFRa. Briefly, PDGFA were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100µl were then transferred to PDGFRa-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-PDGFA pAb, then aspirated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µ stop solution to the wells and read at 450nm immediately. The binding activity of PDGFA and PDGFRa was shown, and this effect was in a dose dependent manner.

Molecular Mass	17.4kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	PDGFA platelet-derived growth factor alpha polypeptide [Homo sapiens]
Official Symbol	PDGFA
Synonyms	PDGFA; platelet-derived growth factor alpha polypeptide; platelet-derived growth factor subunit A; PDGF A chain; PDGF A; PDGF1; platelet derived growth factor alpha chain; PDGF-1; PDGF A-chain; PDGF subunit A; platelet-derived growth factor A chain; platelet-derived growth factor alpha chain; platelet-derived growth factor alpha isoform 2 preproprotein; PDGF-A;
Gene ID	5154

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq [NM_002607](#)

Protein Refseq [NP_002598](#)


MIM [173430](#)

UniProt ID [P04085](#)

Activity



SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA