# Native Mushroom Agaricus bisporus Lectin (ABA/ABL), FITC-Conjugated

**Cat. No.** ABA/ABL-293M **Lot. No.** (See product label)

## SPECIFICATION

| | |
|---|---|
| **Product Overview** | Native Mushroom Agaricus bisporus Lectin (ABA/ABL) - FITC (Fluorescein)<br>Blood Group Specificity: Non-specific<br>Preferred Sugar Specificity: Galactoseβ1-3N-Acetylgalactosamine<br>Inhibiting or Eluting Sugar: Fetuin<br>Divalent Ions: None Required<br>Excitation: 495 nm<br>Emission: 515 nm<br>Mitogenic Activity: No<br>Hazardous Shipping: Non-hazardous<br>Fluorescence: Green |
| **Species** | Mushroom |
| **Source** | Mushroom |
| **Description** | Affinity-pured Agaricus bisporus agglutinin (ABA) is a tetramer composed of two to four isolectins. ABA has two distinct carbohydrate binding sites, one for galactose-β-1,3-N-acetylgalactosamine and another for galactose-β-1,3-N-acetylglucosamine. Agaricus bisporus has an anti-proliferative effect on cancerous cells. ABA can be internalized by clathrin-coated vesicles after binding to surface glycoproteins, thus making ABA an inhibitor of its nuclear import of signal-dependent proteins. Agaricus bisporus Lectin (ABA/ABL) is labeled with fluorescein isothiocyanate (FITC) and has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for thi |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---|---|
| | s lectin. FITC conjugates have been used in a variety of immunohistochemical and flo w cytometry applications. |
| **Form** | Liquid |
| **Stability** | Shelf Life: 1 year |
| **Storage** | At 2-8 centigrade. |
| **Concentration** | 2 mg/mL |

## GENE INFORMATION

| | |
|---|---|
| **Official Symbol** | ABA/ABL |
| **Synonyms** | Agaricus bisporus Lectin; ABA/ABL |

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA