Native Lentil Lens culinaris Lectin (LCA/LCH), TRITC-Conjugated

Cat. No. LCA/LCH-498L **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Native Lentil Lens culinaris Lectin (LCA/LCH) - TRITC (Rhodamine) Blood Group Specificity: Non-specific Preferred Sugar Specificity: α-Mannose, α-Glucose Inhibiting or Eluting Sugar: α-Methylmannoside + α-Methylglucoside Divalent Ions: Ca++, Mn++ Excitation: 545-555 nm Emission: 570-580 nm Mitogenic Activity: Yes Hazardous Shipping: Non-hazardous Fluorescence: Orange
Species	Lentil
Source	Lentil
Description	This lectin is useful in affinity chromatography columns for the separation of glycoconjugates. The two isomers LCA-A and LCA-B agglutinates human red blood cells, although not blood group specific. Activity of LCA is determined by haemagglutination with human blood and agglutinates a 2% suspension of human erythrocytes at a lectin concentration of 8µg/ml. LCA requires calcium and manganese ions for binding. Lens culinaris (Lentil) Lectin (LCA/LCH) is labeled with tetramethylrhodamine isothiocyanate (TRITC) and has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for this lectin.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	TRITC is a bright orange or red-fluorescent dye with excitation ideally suited to the 532-nm laser line. TRITC labeled LCA/LCH can be used for cellular imaging applications.
Form	Liquid
Purity	High Purity Grade
Stability	Shelf Life: 1 year
Storage	At -20 centigrade.
Concentration	5 mg/mL

GENE INFORMATION

Official Symbol	LCA/LCH
Synonyms	Lens culinaris Lectin; LCA/LCH

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA