Native Japanese pagoda tree Styphnolobium japonicum Lectin (SJA)

Cat. No. SJA-717J **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Native Japanese pagoda tree Styphnolobium japonicum Lectin (SJA) - Semi-Pure Preferred Sugar Specificity: β-N-Acetylgalactosamine Inhibiting or Eluting Sugar: N-Acetylgalactosamine Divalent Ions: Ca++, Mn++ Mitogenic Activity: No Hazardous Shipping: Non-hazardous
Species	Japanese pagoda tree
Source	Japanese pagoda tree
Description	Styphnolobium japonicum agglutinin (SJA) is fractionated from Japanese pagoda tree seeds. It consists of two subunits. The subunits can be separated into two subfractions, a D-galactose/N-acetyl-D-galactosamine specific lectin (B-SJA-I) and a D-mannose/D-glucose specific lectin (B-SJA-II). SJA agglutinates all blood group types but has greater affinity for A erythrocytes than of B types, than of O (-SA) types. Binding experiments of the lectin with frozen sections of human kidneys shows specific binding to the endothelia in specimens from blood groups B or AB, thus indicating a D-galactose/N-acetyl-D-galactosamine receptor specificity. SJA lacks mitogenic and immune-suppressive activity.
Form	Powder

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	At -20 centigrade.

GENE INFORMATION

Official Symbol	SJA
Synonyms	Styphnolobium japonicum Lectin; SJA

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA