

Recombinant Human TRAV10 Protein, His-tagged

Cat. No. TRAV10-70H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TRAV10 Protein, fused to His-tag, was expressed in E. coli.
Species	Human
Source	E.coli
Description	T cell receptors recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of antigen presenting cells (APC). Each T cell receptor is a dimer consisting of one alpha and one beta chain or one delta and one gamma chain. In a single cell, the T cell receptor loci are rearranged and expressed in the order delta, gamma, beta, and alpha. If both delta and gamma rearrangements produce functional chains, the cell expresses delta and gamma. If not, the cell proceeds to rearrange the beta and alpha loci. This region represents the germline organization of the T cell receptor alpha and delta loci. Both the alpha and delta loci include V (variable), J (joining), and C (constant) segments and the delta locus also includes diversity (D) segments. The delta locus is situated within the alpha locus, between the alpha V and J segments. During T cell development, the delta chain is synthesized by a recombination event at the DNA level joining a D segment with a J segment; a V segment is then joined to the D-J gene. The alpha chain is synthesized by recombination joining a single V segment with a J segment. For both chains, the C segment is later joined by splicing at the RNA level. Recombination of many different V segments with several J segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random additional of nucleotides by terminal

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

deoxynucleotidyltransferase. Five variable segments can be used in either alpha or delta chains and are described by TRAV/DV symbols. Several V and J segments of the alpha locus are known to be incapable of encoding a protein and are considered pseudogenes.

Form	Liquid. In Tris buffer(20mM Tris,150mM NaCl, pH 9.0).
Molecular Mass	~27.8 kDa
AA Sequence	MNQVEQSPQSLIILEGKNCTLQCNYTVSPFSNLRWYKQDTGRGPVSLTIMTFSENTK SNGRYTATLDADTKQSSLHITASQLSDSASYICVVSDRGSTLGRLYFGRGTQLTVWP DGGGGSGGGGSGGGGSGGGGSEADYQTPRYLVIGTGKKITLECSQTMGHDKMY WYQQDPGMELHLIHYSYGVNSTEKGDLSSSESTVSRIRTEHFPLTLESARPSHTSQYL CASSENIGTAYEQYFGPGTRTLVTEHHHHHH
Purity	>90%
Storage	Short Term Storage at 4 centigrade, Long Term, please prepare aliquots with 20% glycerol and store at -20-80 centigrade. Avoid freeze/thaw cycles.
Concentration	0.1 mg/mL

GENE INFORMATION

Gene Name	TRAV10 T cell receptor alpha variable 10 [Homo sapiens (human)]
Official Symbol	TRAV10
Synonyms	TCRAV10S1; TCRAV24S1
Gene ID	28676

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID


A0A0B4J240

SDS-PAGE



15% SDS-PAGE analysis profiles of purified StcrV2, His-tagged (CBB stained).

WB



Western blotting analysis profiles of purified StcrV2, His-tagged (Antibody Rat Anti-

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA



his (HRP)).

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA