Recombinant Full Length Rat C-C Chemokine Receptor Type 2(Ccr2) Protein, His-Tagged

Cat. No. RFL19268RF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Rat C-C chemokine receptor type 2(Ccr2) Protein (O55193) (1-373aa), fused to N-terminal His tag, was expressed in E. coli.
Species	Rat
Source	E.coli
ProteinLength	Full Length (1-373)
Form	Lyophilized powder
AA Sequence	MEDSNMLPQFIHGILSTSHSLFPRSIQELDEGATTPYDYDDGEPCHKTSVKQIGAWILPP LYSLVFIFGFVGNMLVIIILISCKKLKSMTDIYLFNLAISDLLFLLTLPFWAHYAANEWV FGNIMCKLFTGLYHIGYFGGIFFIILLTIDRYLAIVHAVFALKARTVTFGVITSVVTWVV AVFASLPGIIFTKSEQEDDQHTCGPYFPTIWKNFQTIMRNILSLILPLLVMVICYSGILH TLFRCRNEKKRHRAVRLIFAIMIVYFLFWTPYNIVLFLTTFQEFLGMSNCVVDMHLDQ AM QVTETLGMTHCCVNPIIYAFVGEKFRRYLSIFFRKHIAKNLCKQCPVFYRETADR VSSTF TPSTGEQEVSVGL
Purity	Greater than 90% as determined by SDS-PAGE.
Applications	SDS-PAGE
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	for up to one week.
Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.
Storage Buffer	Tris/PBS-based buffer, 6% Trehalose, pH 8.0
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20℃/-80℃. Our default final concentration of glycerol is 50%. Customers could use it as reference.

GENE INFORMATION

Gene Name	Ccr2
Synonyms	Ccr2; Cmkbr2; C-C chemokine receptor type 2; C-C CKR-2; CC-CKR-2; CCR-2; CCR2; CD antigen CD192
UniProt ID	O55193

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA