

Recombinant Human CD3E & CD3D, His & Flag tag "

Cat. No. CD3E&D-308H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	A DNA sequence encoding the extracellular domain (Met1-Ala105) of human CD3D (P04234) was fused with a flag tag at the C-terminus, constructed the plasmid 1; A DNA sequence encoding the extracellular domain (Met1-Asp126) of human CD3E (P07766) was fused with a polyhistidine tag at the C-terminus, constructed the plasmid 2. The two plasmids were co-expressed and the human CD3D/CD3E heterodimer was purified.
Source	Human Cells
ProteinLength	1-105 a.a.
Description	T-cell surface glycoprotein CD3 delta/epsilon chain, also known as CD3D and CD3E respectively, are single-pass type I membrane proteins. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. CD3D, together with CD3- epsilon(CD3E) , CD3-gamma and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). The immunofluorescence defined that CD3 epsilon(CD3E0) was strongly stained bound to either CD 3 gamma or CD 3 delta. The recognition by this panel of anti-CD3 antibodies of CD3-gamma/ epsilon and CD3-delta/ epsilon complexes and not of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

CD3-epsilon alone was assessed by immunoprecipitation.

Molecular Mass

The recombinant heterodimer of human CD3D&CD3E comprises 272 (126+146) amino acids and has a calculated molecular mass of 31.2 (14.4+16.8)KDa. As a result of glycosylation, the apparent molecular mass of human CD3D&CD3E heterodimer is approximately 20, 22 and 26 KDa in SDS-PAGE under reducing conditions.

Predicted N terminal

Phe 22 & Asp 23

Purity

(43.8+13.1+41.9) % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Formulation

Lyophilized from sterile PBS, pH 7.4.

Storage

Store it under sterile conditions at -70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

A hardcopy of COA with reconstitution instruction is sent along with the products.

GENE INFORMATION

Synonyms

CD3e molecule, epsilon (CD3-TCR complex); T3E; TCRE; FLJ18683; CD3E; T-cell surface glycoprotein CD3 epsilon chain; CD3-epsilon; T-cell surface antigen T3/Leu-4 epsilon chain; CD3e antigen, epsilon polypeptide (TiT3 complex); T-cell antigen receptor complex, epsilon subunit of T3; CD3E antigen, epsilon polypeptide; CD3e antigen; CD3d molecule, delta (CD3-TCR complex); T3D; CD3-DELTA; CD3D; T-cell surface glycoprotein CD3 delta chain; OKT3, delta chain; CD3 antigen, delta subunit; T-cell receptor T3 delta chain; CD3d antigen, delta polypeptide (TiT3 complex); CD3d antigen

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA