

Recombinant Full Length Human C-X-C Chemokine Receptor Type 2(Cxcr2) Protein, His-Tagged

Cat. No. RFL26660HF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Human C-X-C chemokine receptor type 2(CXCR2) Protein (P25025) (1-360aa), fused to N-terminal His tag, was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	Full Length (1-360)
Form	Lyophilized powder
AA Sequence	MEDFNMESDSFEDFWKGEDLSNYSYSTLPPFLD AAPCEPESLEINKYFVVIYALV FL LSLGNSLVMLVILYSRVGRSVTDVYLLNLALADLLFALTLP IWAASKVNGWIFGTF LCK VVSL LKEVNFYSGILLACISVD RYLAIVHATRTL TQKRYLVKFICLSIWGLSLLLA LPV LLFRRTVYSSNVSPACYEDMGNNTANWRMLLRILPQSFGFIVPLIMLFCYGF LRTLFK AHMGQKHRAMRVIFAVVLIFLLCWLPYNLVLLADTL MRTQVIQETCERNHI DRALDATE ILGILHSCLNPLIYAFIGQKFRHGLLKILAIHGLISKDSL PKDSRPSFVGSS SGHTSTTL
Purity	Greater than 90% as determined by SDS-PAGE.
Applications	SDS-PAGE
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	for up to one week.
Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.
Storage Buffer	Tris/PBS-based buffer, 6% Trehalose, pH 8.0
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
GENE INFORMATION	
Gene Name	CXCR2
Synonyms	C-X-C chemokine receptor type 2; CD 182; CD182; CD182 antigen; CDw128b; Chemokine (CXC) receptor 2; CMKAR2; CXC-R2; CXCR 2; CXCR-2; CXCR2; CXCR2_HUMAN; GRO/MGSA receptor; High affinity interleukin-8 receptor B; IL 8 receptor type 2; IL 8R B; IL-8 receptor
UniProt ID	P25025