

Active Recombinant Human TGFB2, Animal Free

Cat. No. TGFB2-127H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human TGF-beta2 is a 27.08 kDa protein composed of two identical 118 amino acid polypeptide chains linked by a single disulphide bond. Human recombinant protein expressed in Nicotiana benthamiana. Recombinant human TGF-beta-2 contains a 6-His-tag at the N-terminal end, is produced by transient expression in non-transgenic plants and is purified by sequential chromatography (FPLC). This product contains no animal-derived components or impurities. Animal free product.
-------------------------	--

Species	Human
----------------	-------

Source	Nicotiana Benthamiana
---------------	-----------------------

Description	Recombinant human TGF-beta2 is a 27.08 kDa protein composed of two identical 118 amino acid peptide chains linked by a single disulphide bond. Transforming growth factor-beta is a family of five related cytokines that have been shown on a wide variety of normal and neoplastic cells, indicating the importance of these homo-dimer proteins as multi-functional regulators of cellular activity. The three mammalian isoforms of TGF-beta (TGF-beta1, TGF-beta2 and TGF-beta3) signal through the same receptor and elicit similar biological responses. They are involved in physiological processes as embryogenesis, tissue remodelling and wound healing.
--------------------	--

Form	Lyophilized from a Tris HCl 0.05M buffer at pH 7.4.
-------------	---

Bio-activity	The biological activity of TGF-beta2 is measured in culture by its ability to inhibit the mink lung epithelial (Mv1Lu) cells proliferation. ED50 ≤ 50ng/ml
---------------------	--

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Recombinant human TGF-beta2 is a 27.08 kDa protein composed of two identical 118 amino acid polypeptide chains linked by a single disulphide bond.
AA Sequence	HHHHHHALDAAYCFRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCAGACP YLWSSDTQHRSRVLSLYNTIN PEASASPCCVSQDLEPLTILYYIGKTPKIEQLSNMIVK SCKCS
Endotoxin	< 0.04="" eu="" ug="" protein="" (lal="">
Purity	>97% by SDS-PAGE gel
Applications	Cell culture, Western blot
Storage	This lyophilized preparation is stable at 2-8o C for short term, long storage it should be kept at -20oC. Reconstituted protein should be stored in working aliquots at –20°C. It is recommended to add a carrier protein (0.1% HSA or BSA). Repeated freezing and thawing is not recommended.
Reconstitution	Lyophilized protein should be reconstituted in water to a concentration of 25-50 ng / ul. Due to the protein nature, dimmers and multimers may be observed. Upon reconstitution, It can be stored in working aliquots at –20°C for future use. Optimal reconstitution please follow batch Quality Control sheet instructions.

GENE INFORMATION

Gene Name	TGFB2 transforming growth factor, beta 2 [Homo sapiens]
Official Symbol	TGFB2
Synonyms	TGFB2; transforming growth factor, beta 2; transforming growth factor beta-2; G-TSF; cetermin; polyergin; BSC-1 cell growth inhibitor; glioblastoma-derived T-cell

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	suppressor factor; TGF-beta2; MGC116892;
Gene ID	7042
mRNA Refseq	NM_001135599
Protein Refseq	NP_001129071
MIM	190220
UniProt ID	P61812
Chromosome Location	1q41
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Cell cycle, organism-specific biosystem; Cell cycle, conserved biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem;
Function	beta-amyloid binding; cytokine activity; growth factor activity; protein binding; contributes_to protein binding; protein heterodimerization activity; protein homodimerization activity; receptor binding; receptor signaling protein serine/threonine kinase activity; transforming growth factor beta receptor binding; type II transforming growth factor beta receptor binding; type II transforming growth factor beta receptor binding;