

Recombinant Full Length Escherichia Coli Cdp-Diacylglycerol-Glycerol-3-Phosphate 3-Phosphatidyltransferase(Pgsa) Protein, His-Tagged

Cat. No. RFL3921EF Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Escherichia coli CDP-diacylglycerol--glycerol-3-phosphate 3-phosphatidyltransferase(pgsA) Protein (P0ABF8) (2-182aa), fused to N-terminal His tag, was expressed in E. coli.
Species	E.coli
Source	E.coli
ProteinLength	Full Length of Mature Protein (2-182)
Form	Lyophilized powder
AA Sequence	QFNIPTLLTLFRVILIPFFVLVFYLPVTWSPFAAALIFCVAAVTDWFDGFLARRWNQST R FGAFLDPVADKVLVAIAMVLVTEHYHSWWVTLPAAATMIAREIIISALREWMAELGK RSSV AVSWIGKVKTAAQMVALAWLLWRPNIWVEYAGIALFFVAAVLTLSMLQYLS AARADLLD Q
Purity	Greater than 90% as determined by SDS-PAGE.
Applications	SDS-PAGE
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.
Storage Buffer	Tris/PBS-based buffer, 6% Trehalose, pH 8.0
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
GENE INFORMATION	
Gene Name	pgsA
Synonyms	pgsA; b1912; JW1897; CDP-diacylglycerol--glycerol-3-phosphate 3-phosphatidyltransferase; Phosphatidylglycerophosphate synthase; PGP synthase
UniProt ID	P0ABF8