

Recombinant Full Length Nicotiana Tabacum Cytochrome B-C1 Complex Subunit Rieske-4, Mitochondrial Protein, His-Tagged

Cat. No. RFL12144NF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Nicotiana tabacum Cytochrome b-c1 complex subunit Rieske-4, mitochondrial Protein (P51134) (25-236aa), fused to N-terminal His tag, was expressed in E. coli.
Species	Nicotiana tabacum
Source	E.coli
ProteinLength	Full Length of Mature Protein (25-236)
Form	Lyophilized powder
AA Sequence	SSNSVSHAHDMGLVPDLPPTVAAIKNPTSKIVYDEHNHERYPPGDPSKRAFAYFVLTTGGR FVYASLVRLILKFVLSMSASKDVLALASLEVDLSSIEPGTTVTVKWRGKPVFIRRRRTED DINLANSVDLGSLRDPQQDAERVKSPEWLVVIGVCTHLGCIPLPNAGDFGGWFCPCHGSH YDISGRIRKGPAPYNLEVPTYSFLEENKLLIG
Purity	Greater than 90% as determined by SDS-PAGE.
Applications	SDS-PAGE
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.
Storage Buffer	Tris/PBS-based buffer, 6% Trehalose, pH 8.0
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
GENE INFORMATION	
Gene Name	Nicotiana tabacum Cytochrome b-c1 complex subunit Rieske-4, mitochondrial
Synonyms	Cytochrome b-c1 complex subunit Rieske-4, mitochondrial; Complex III subunit 5-4; Rieske iron-sulfur protein 4; RISP4; Ubiquinol-cytochrome c reductase iron-sulfur subunit 4
UniProt ID	P51134