

Recombinant Human P4HB, MYC/DDK-tagged

Cat. No. P4HB-116H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human P4HB fused with C-terminal MYC/DDK, was expressed in HEK293 Cells.
Species	Human
Source	HEK293
Description	This gene encodes the beta subunit of prolyl 4-hydroxylase, a highly abundant multifunctional enzyme that belongs to the protein disulfide isomerase family. When present as a tetramer consisting of two alpha and two beta subunits, this enzyme is involved in hydroxylation of prolyl residues in procollagen. This enzyme is also a disulfide isomerase containing two thioredoxin domains that catalyze the formation, breakage and rearrangement of disulfide bonds. Other known functions include its ability to act as a chaperone that inhibits aggregation of misfolded proteins in a concentration-dependent manner, its ability to bind thyroid hormone, its role in both the influx and efflux of S-nitrosothiol-bound nitric oxide, and its function as a subunit of the microsomal triglyceride transfer protein complex.
Molecular Mass	55.2 kDa
Purity	>80% as determined by SDS-PAGE and Coomassie blue staining
Storage	Store at -80C. Avoid repeated freeze-thaw cycles. Stable for at least 3 months from receipt of products under proper storage and handling conditions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Concentration	>50 ug/mL as determined by microplate BCA method
Storage Buffer	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.
GENE INFORMATION	
Gene Name	P4HB prolyl 4-hydroxylase, beta polypeptide [Homo sapiens (human)]
Official Symbol	P4HB
Synonyms	P4HB; DSI; GIT; PDI; PHDB; PDIA1; PO4DB; PO4HB; PROHB; ERBA2L; P4Hbeta; prolyl 4-hydroxylase, beta polypeptide; protein disulfide-isomerase; p55; procollagen hydroxylase; prolyl 4-hydroxylase subunit beta; collagen prolyl 4-hydroxylase beta; thyroid hormone-binding protein p55; glutathione-insulin transhydrogenase; cellular thyroid hormone-binding protein; protein disulfide isomerase-associated 1; protein disulfide isomerase/oxidoreductase; protein disulfide isomerase family A, member 1; procollagen-proline, 2-oxoglutarate 4-dioxygenase (proline 4-hydroxylase), beta polypeptide; EC 5.3.4.1
Gene ID	5034
mRNA Refseq	NM_000918
Protein Refseq	NP_000909
MIM	176790
UniProt ID	P07237
Chromosome Location	17q25

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Chylomicron-mediated lipid transport; Collagen biosynthesis and modifying enzymes; Collagen formation
Function	electron carrier activity; procollagen-proline 4-dioxygenase activity; protein binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA