

Recombinant Human MGAT2, His-tagged

Cat. No. MGAT2-5118H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human MGAT2 protein, fused to His-tag at N-terminus, was expressed in E.coli.
Species	Human
Source	E.coli
Description	MGAT2 is a golgi enzyme catalyzing an essential step in the conversion of oligomannose to complex N-glycans. The enzyme has the typical glycosyltransferase domains: a short N-terminal cytoplasmic domain, a hydrophobic non-cleavable signal-anchor domain, and a C-terminal catalytic domain. Mutations in this gene may lead to carbohydrate-deficient glycoprotein syndrome, type II. Transcript variants with a spliced 5' UTR may exist, but their biological validity has not been determined.
Form	Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol
Molecular Mass	50 kDa (439aa)
AA Sequence	MGSSHHHHHH SSGLVPRGSH MRQRKNEALA PPLDAEPAR GAGGRGGDHP SVAVGIRRVSV NVSAASLVPA VPQPEADNLT LRYRSLVYQL NFDQTLRNVD KAGTWAPREL VLVVQVHNRPEYLRLLLLDSL RKAQGIDNVL VIFSHDFWST EINQLIAGVN FCPVLQVFFP FSIQLYPNEF PGSDPRDCPR DLPKNAALKL GCINAEYPDS FGHYREAKFS QTKHHWWWKL HFVWERVKIL RDYAGLILFL EEDHYLAPDF YHVFKKMWKL KQQECPECDV LSLGTYSASR SFYGMADKVD VKTKWKSTEHN MGLALTRNAY QKLIECTDTF CTYDDYNWDW TLQYLTVSCL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	PKFWKVLVPQ IPRIFHAGDC GMHHKKTCP STQSAQIESL LNNNKQYMFP ETLTISEKFT VVAISPPRKN GGWGDIRDHE LCKSYRRLQ
Purity	>85% by SDS - PAGE
Applications	SDS-PAGE
Storage	Can be stored at +4centigrade short term (1-2 weeks). For long term storage, aliquot and store at -20centigrade or -70centigrade. Avoid repeated freezing and thawing cycles.
Concentration	1.0 mg/ml (determined by Bradford assay)
GENE INFORMATION	
Gene Name	MGAT2 mannosyl (alpha-1,6-)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase [Homo sapiens (human)]
Official Symbol	MGAT2
Synonyms	MGAT2;GNT2; CDG2A; CDGS2; GNT-II; GLCNACTII;alpha-1,6-mannosyl-glycoprotein 2-beta-N-acetylglucosaminyltransferase;glcNAc-T II;mannoside acetylglucosaminyltransferase 2;Beta-1,2-N-acetylglucosaminyltransferase II;N-glycosyl-oligosaccharide-glycoprotein N-
Gene ID	4247
mRNA Refseq	NM_002408.3
Protein Refseq	NP_002399.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	602616
UniProt ID	Q10469
Chromosome Location	14q21
Pathway	Asparagine N-linked glycosylation;Metabolism of proteins;N-Glycan biosynthesis;N-glycan antennae elongation in the medial/trans-Golgi;N-glycan biosynthesis, complex type
Function	alpha-1,6-mannosylglycoprotein 2-beta-N-acetylglucosaminyltransferase activity;carbohydrate binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA